

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 36, 2026, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 36, 2026, 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Ana Fortič, Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, James Pierson (USA), Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednici/Redattrici/Editors:

Martina Orlando-Bonaca, Ana Fortič

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*[®]
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment* / *Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*[®]

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano* / *National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2026.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base* / *Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza* / *This work is licensed under a Creative Commons BY 4.0.*



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / *INDICE GENERALE* / *CONTENTS*

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Alen SOLDÓ, Rigers BAKIU & Sherif DURMISHAJ
The First Record of Bastard Grunt *Pomadasys incisus* (Bowdich, 1825) in Albanian Waters (Adriatic Sea) 1
Prvi zapis o pojavljanju neprave prašičevke Pomadasys incisus (Bowdich, 1825) v albanskih vodah (Jadransko morje)

Pero UGARKOVIĆ, Ilija ČETKOVIĆ, Olivera MARKOVIĆ, Aleksandar JOKSIMOVIĆ, Nikola ĐORĐEVIĆ & Jakov DULČIĆ Additional Records of the Bastard Grunt, *Pomadasys incisus* (Bowdich, 1825), from the Eastern Adriatic Sea 7
Dodatne najdbe neprave prašičevke, Pomadasys incisus (Bowdich, 1825), iz vzhodnega Jadranskega morja

Rüştü KIRMAN & Murat BİLECENOĞLU Significant Range Expansion of *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) in the Aegean Sea Based on Scuba and Photographic Observations 15
Znatno razširjanje areala vrste Sepioteuthis lessoniana (Cephalopoda: Loliginidae), v Egejskem morju na podlagi podvodnih opazovanj in fotografiranja

Adrian BRAJKOVIĆ, Iris MATULJA & Neven IVEŠA
Conquering the North: New Occurrence of the Common Lionfish, *Pterois miles* (Bennett, 1828), in the Adriatic Sea 21
Osvajanje severa: novi zapis o pojavljanju navadne plamenke, Pterois miles (Bennett, 1828), v Jadranskem morju

Alan DEIDUN, Bruno ZAVA, Maria CORSINI-FOKA, Arnold SCIBERRAS & Alessio MARRONE First Record of the Devil Firefish *Pterois miles* (Bennett, 1828) (Actinopterygii: Scorpaenidae) from Maltese Coastal Waters 29
Prvi zapis o pojavljanju plamenke Pterois miles (Bennett, 1828) (Actinopterygii: Scorpaenidae) v malteških obalnih vodah

Chirine HUSSEIN, Amir IBRAHIM, Firas ALSHAWY, Mouina BADRAN & Rahaf ABO ASA First Mediterranean Record of the Diagonal Butterflyfish, *Chaetodon fasciatus* Forsskal, 1775, Reported from Syrian Waters 35
Prvi sredozemski zapis o pojavljanju poševnoprogame ščetinozobke, Chaetodon fasciatus Forsskal, 1775, iz sirskih voda

Houssein ELBARAASI, Tarek SHOEIB, Mona SAID & Laith A. JAWAD A Further Record of the Barred Knifejaw, *Oplegnathus fasciatus* (Temminck & Schlegel, 1844), a Pacific Species, from the Mediterranean: A New Record from Benghazi, Libya 41
Novi zapis o vrsti Oplegnathus fasciatus (Temminck & Schlegel, 1844), pacifiški vrsti, v Sredozemlju: nova najdba iz Bengazija v Libiji

Alen SOLDÓ & Rigers BAKIU Rapid Increase in Records of the Invasive Silver-cheeked Toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Adriatic Sea 49
Hitro naraščanje pojavov invazivne srebrnoproge napihvalke Lagocephalus sceleratus (Gmelin, 1789) v Jadranskem morju

Gerasimos KONDYLATOS, Konstantinos KALAENTZIS, Styliani MINOUDI & Maria CORSINI-FOKA Further Molecular Identification Confirms the Occurrence of *Lagocephalus guentheri* Miranda Ribeiro, 1915 in the Aegean Coastal Waters of Greece 55
Nadaljnja molekularna identifikacija potrjuje prisotnost vrste Lagocephalus guentheri Miranda Ribeiro, 1915 v egejskih obalnih vodah Grčije

Deniz ERGÜDEN & Cem ÇEVİK Length–Weight and Length–Length Relationships, and Condition Factor of *Ambassis dussumieri* Cuvier, 1828, in the Northeastern Mediterranean, Türkiye 65
Dolžinsko-masni odnos med dolžino in kondicijskim faktorjem pri vrsti Ambassis dussumieri Cuvier, 1828, v severovzhodnem Sredozemskem morju, Turčija

Okan AKYOL & HALİL ŞEN Unexpected Occurrence of Prussian Carp *Carassius gibelio* (Cyprinidae) in Homa Lagoon (Izmir Bay, Aegean Sea) 73
Nepričakovani pojav srebrnega koreslja Carassius gibelio (Cyprinidae) v laguni Homa (Izmirski zaliv, Egejsko morje)

SREDOZEMSKÉ HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
MEDITERRANEAN SHARKS AND RAYS

Terry CARBON, Emily GIGNON, Justine LALLAU-VAZZOLER, Hugo MENARD, Claudio BARRIA, Ana I. COLMENERO & Nicolas ZIANI Precopulatory Behaviour of *Pteroplatytrygon violacea* (Myliobatiformes: Dasyatidae) in the Northwestern Mediterranean 81
Predkopulacijsko vedenje vijoličnega morskega biča (Pteroplatytrygon violacea) (Myliobatiformes: Dasyatidae) v severozahodnem Sredozemlju

Hakan KABASAKAL Distribution of <i>Odontaspis ferox</i> in the Mediterranean Sea: Insights from Spatial and Temporal Analyses 87 <i>Razširjenost vrste Odontaspis ferox v Sredozemskem morju: Spoznanja na podlagi prostorskih in časovnih analiz</i>	
Hristina GELEVSKA, Borut MAVRIČ, Lovrenc LIPEJ & Christian CAPAPÉ Is the Gulf of Trieste a potential nursery area for some elasmobranch species? 99 <i>Ali je Tržaški zaliv potencialno območje odraščanja (jaslice) za določene vrste hrustančnic?</i>	
IHTIOFAVNA ITTIOFAUNA ICHTHYOFAUNA	
Cem DALYAN, Yunus GÖNÜL, Mahmud Samed ŞAHİNOĞULLARI & Hakan KABASAKAL New data on the Occurrence and Morphology of the Armless Snake Eel, <i>Dalophis imberbis</i> (Ophichthidae), from the Northeastern Mediterranean Sea 119 <i>Novi podatki o pojavljanju in morfologiji kačaste jegulje, Dalophis imberbis (Ophichthidae), iz severovzhodnega Sredozemskega morja</i>	
Chirine HUSSEIN, Firas ALSHAWY & Amir IBRAHIM First Record of the Mediterranean Dealfish, <i>Trachipterus trachipterus</i> (Gmelin, 1789) (Trachipteridae), in Syrian Marine Waters 127 <i>Prvi zapis o pojavljanju kosice, Trachipterus trachipterus (Gmelin, 1789) (Trachipteridae), v sirskih morskih vodah</i>	
Igor AGOSTINI & Okan AKYOL On the Occurrence of the Currently Largest Recorded <i>Pagellus erythrinus</i> (Sparidae) in the Tyrrhenian Sea (Italy) 133 <i>O pojavljanju trenutno največjega evidentiranega primerka vrste Pagellus erythrinus (Sparidae) v Tirenskem morju (Italija)</i>	
Ismail Burak DABAN & Yusuf ŞEN Substantiated Record and Preliminary Biological Insights of <i>Microlophrys dalmatinus</i> (Steindachner & Kolombatović, 1883) from the Sea of Marmara, Türkiye 141 <i>Potrjen zapis o pojavljanju in predhodna biološka spoznanja o vrsti Microlophrys dalmatinus (Steindachner & Kolombatović, 1883) iz Marmarskega morja, Turčija</i>	
Zeliha ERDOĞAN, Gülçin ULUNEHİR AYDIN, Hatice TORCU-KOÇ & Tuğba ANBAROĞLU Population Study of <i>Trachurus mediterraneus</i> Focused on Reproductive Biology in Edremit Bay, Northern Aegean Sea 151 <i>Razmnoževalna biologija sredozemskega šura (Trachurus mediterraneus) v sklopu populacijske raziskave v Edremitnem zalivu (severno Egejsko morje)</i>	
MORSKA FAVNA FAUNA MARINA MARINE FAUNA	
Francesco TIRALONGO, Paola LEOTTA & Riccardo MARTELLUCCI <i>Physalia physalis</i> in the central Mediterranean Sea: Recent observations associated with mass strandings of <i>Velella velella</i> and surface circulation dynamics 165 <i>Portugalska ladjica (Physalia physalis) v osrednjem Sredozemskem morju: nedavna opažanja v povezavi z masovnimi nasedanji morskega jadrčka (Velella velella) in dinamiko površinskega kroženja vode</i>	
Andrea LOMBARDO, Juba CHABANE & Rachida GHALMI First Contribution to the Study of the "Sea Slug" Fauna (Gastropoda, Heterobranchia) of the Algerian coast 177 <i>Prvi prispevek k proučevanju favne „goliš morskih polžev“ (Gastropoda, Heterobranchia) alžirske obale</i>	
Borut MAVRIČ, Tjaša PREMRL, Martin MAVRIČ, Tina MIRT, Neža LEBAN, Tihomir MAKOVEC & Lovrenc LIPEJ Assessing Soft-Bottom Epibenthic Communities: Methodological Insights into Dredging and Video Surveys 203 <i>Ocenjevanje epibentoških združb mehkega dna: metodološki vpogledi v dredžanje in video pregleda</i>	
FAVNA FAUNA FAUNA	
Kim LEBAN, Iztok ŠKORNIK, Špela ČONČ, Mateja BREG VALJAVEC, Žan KURALT, Lenart ŠTAUT, Jure TIČAR, Katarina POLAJNAR HORVAT, Primož GAŠPERIČ, Aleš SMREKAR & Melita VAMBERGER Habitat Insights for the European Pond Turtle (<i>Emys orbicularis</i>) in the Coastal Wetland Area Sečovlje Salina Nature Park, Slovenia 221 <i>Habitatne značilnosti močvirske sklednice (Emys orbicularis) v krajinskem parku Sečoveljske soline, Slovenija</i>	
Leon CIMERMAN & Matija KRIŽNAR Najdbe pleistocenske avifavne v kamnolomu Črni Kal (Primorska, Slovenija) 243 <i>Pleistocene Avifauna Finds in the Črni Kal Quarry (Primorska Region, Slovenia)</i>	
Kazalo k slikam na ovitku 255 Index to images on the cover 255	

received: 2026-03-30

DOI 10.19233/ASHN.2026.11

UNEXPECTED OCCURRENCE OF PRUSSIAN CARP *CARASSIUS GIBELIO* (CYPRINIDAE) IN HOMA LAGOON (IZMIR BAY, AEGEAN SEA)

*Okan AKYOL & Halil ŞEN*Ege University Faculty of Fisheries, 35440 Urla, Izmir, Türkiye
e-mail: okan.akyol@ege.edu.tr

ABSTRACT

On 24 February 2026, a female specimen of *Carassius gibelio* was caught using a trammel net targeting grey mullets in Homa Lagoon, located on the coast of Izmir Bay, at a depth of 1 m. Homa is a highly saline environment, with salinity values ranging from 35.6 ‰ to 63.7 ‰. Therefore, the occurrence of *C. gibelio*, a potamodromous fish species, in its waters represents an unexpected phenomenon. This paper not only presents a remarkable occurrence of this invasive species in an unusual habitat, but also provides an additional contribution to the fish fauna records of Homa Lagoon, Izmir Bay.

Key words: Salty lagoon, unusual habitat, measurement, Mediterranean

OCCORRENZA INASPETTATA DELLA CARPA DI PRUSSIA *CARASSIUS GIBELIO* (CYPRINIDAE) NELLA LAGUNA DI HOMA (BAIA DI IZMIR, MAR EGEO)

SINTESI

Il 24 febbraio 2026, un esemplare femmina di *Carassius gibelio* è stato catturato mediante una rete da imbrocco utilizzata per la pesca dei cefali grigi nella laguna di Homa, situata sulla costa della Baia di Izmir, a una profondità di 1 m. Homa è un ambiente altamente salino, con valori di salinità compresi tra 35,6 ‰ e 63,7 ‰. Pertanto, la presenza di *C. gibelio*, una specie ittica potamodroma, nelle sue acque rappresenta un fenomeno inatteso. Questo lavoro non solo documenta una notevole presenza di questa specie invasiva in un habitat insolito, ma fornisce anche un ulteriore contributo ai dati sulla fauna ittica della laguna di Homa, nella Baia di Izmir.

Parole chiave: laguna salata, habitat insolito, misurazione, Mediterraneo

INTRODUCTION

Cypriniformes are among the most evolutionarily successful freshwater fish, characterized by wide distribution, high abundance, and remarkable species diversity. This paper focuses on the Prussian carp *Carassius gibelio* (Bloch, 1782), as this species still retains a strong ability to colonize new areas and is well known for invading diverse water bodies (Afanasyev *et al.*, 2025). Nowadays, *C. gibelio* has attracted considerable interest from aquatic biologists. It is an invasive species with a widespread distribution resulting from introductions and translocations across most European countries (Perdikaris *et al.*, 2012).

Carassius gibelio inhabits a wide variety of still water bodies and lowland rivers usually associated with submerged vegetation or periodic flooding. It displays strong tolerance to low oxygen concentrations and pollution. Its diet includes plankton, benthic invertebrates, plant material, and detritus, *C. gibelio* spawns on submerged vegetation in shallow, warm shores and is capable of developing from unfertilized eggs (gynogenesis). Its life span reaches up to about 10 years. Length at maturity

is 10.3 cm, the maximum total length is 46.6 cm (common total length: 20.0 cm), and the maximum published weight is 3.0 kg (Froese & Pauly, 2025).

Carassius gibelio is distributed across Europe and Asia. Clear and definite data on its native range in Europe are lacking due to historical introductions, confusion with *Carassius auratus*, and complex breeding patterns. At present, the species is widely distributed and commonly stocked together with *Cyprinus carpio*, which is transported throughout Europe. It is absent from the northern Baltic basin, Iceland, Ireland, Scotland, and Mediterranean islands (Froese & Pauly, 2025). The species has minimal commercial value, which makes it an undesirable target for fisheries (Perdikaris *et al.*, 2012).

The Prussian carp is listed by the IUCN as one of the 100 worst invasive species in the world. As vector, they were introduced to Türkiye from Europe for fish release purposes. Following its introduction, *C. gibelio* spread rapidly and uncontrollably to other lakes (Uysal & Boz, 2018). This study presents unexpected field data on this species' appearance in a new region, specifically, its first observation in a high-salinity lagoon basin in Izmir Bay, Aegean Sea.



Fig. 1: *Carassius gibelio* caught from Homa Lagoon (ESFM-PIS/2026-01) with a black peritoneum. Scale bar: 50 mm.

Sl. 1: *Carassius gibelio* (srebrni koreselj), ujet v laguni Homa (ESFM-PIS/2026-01), s črnim peritonejem (trebušno mreno). Merilo: 50 mm.




Fig. 2: Sampling site (red dot) of *Carassius gibelio* in Homa Lagoon, Izmir Bay, Aegean Sea.

Sl. 2: Lokacija vzorčenja (rdeča pika) za vrsto *Carassius gibelio* v laguni Homa, Izmirski zaliv, Egejsko morje.

MATERIAL AND METHODS

On 24 February 2026, a female specimen of *Carassius gibelio* (Fig. 1) was caught using a trammel net targeting grey mullets in Homa Lagoon, located on the coast of Izmir Bay (38°32'00" N, 26°52'51" E) at a depth of 1 m (Fig. 2).

Homa Lagoon, situated near the Gediz River delta in the northeastern part of the Izmir Bay, is an important wetland covering an area of approximately 1824 hectares and serving as a feeding, growth, and breeding area for many fish and bird species. Since 1986, the lagoon has been used as an experimental fishing area by the Faculty of Fisheries of Ege University. It has also been designated as a wildlife conservation area since 1984 and has been protected under the Ramsar Convention since 1998 (Akyol, 2005). A total of 39 fish species have been identified within the lagoon to date (Alpbaz & Kınacıgil, 1988; Acarlı *et al.*, 2009). The sample fish was fixed in 6% formaldehyde solution and preserved in the fish collection of the Faculty of Fisheries, Ege University (ESFM-PIS/2026-01). Terminology followed Soto *et al.* (2024), who evaluated the various terms used in invasion science and proposed a simplified and standardized nomenclature.

RESULTS AND DISCUSSION

The specimen was identified as a *Carassius gibelio* based on the following characters: body silvery-brown, peritoneum black; last simple anal and dorsal-fin rays strongly serrated; 29 scales along the lateral line (Fig. 1). Selected morphometric measurements and proportions expressed as percentages of total length (TL%) are presented in Table 1.

Carassius gibelio was first reported in Türkiye from Lake Gala in the 1980s (Baran & Ongan, 1988). Subsequently, the species spread to almost all inland waters of Thrace and Anatolia (Ekmekçi *et al.*, 2013). Since then, the Prussian carp has probably become the most dominant invasive non-native fish species and is now considered naturalized in more than 230 lakes, rivers, streams, ponds, and reservoirs throughout Türkiye (Ekmekçi *et al.*, 2013; Yerli *et al.*, 2014). The dominance of *C. gibelio* is attributed to its broad environmental tolerance and flexible adaptive biological strategies, including omnivorous feeding habits, exploitation of variable habitats (slow-running lotic, lentic, and transitional systems with low salinity), intense reproductive activity, and active competition for food, spawning sites, and spawning substrates (Pardikaris *et al.*, 2012). It is

Tab. 1: Morphometric measurements and proportions expressed as percentages of total length (TL%) for *Carassius gibelio* captured in Homa Lagoon, Izmir Bay, Aegean Sea.

Tab. 1: Morfometrične meritve in deleži, izraženi kot odstotki celotne dolžine (TL %), za primerek vrste *Carassius gibelio*, ujetega v laguni Homa, Izmirski zaliv, Egejsko morje.

Morphometrics	mm	TL%
Total length (TL)	255	-
Standard length (SL)	210	82.4
Head length (HL)	55	21.6
Eye diameter	11	4.3
Snout length	15	5.9
Body depth	81	31.8
Predorsal length	94	36.9
Preanal length	155	60.8
Prepectoral length	52	20.4
Meristic counts		
Dorsal fin	III + 17	
Anal fin	III + 5	
Pectoral fin	18	
Ventral fin	I + 8	
Lateral line	29	
Weight (g)	305	

therefore not surprising for the highly adaptable Prussian carps to occur in some Turkish brackish lagoons, such as Liman or Karaboğaz, Akyatan, and Peso lagoons (Ekmekçi *et al.*, 2013). These lagoons are characterized by slightly saline (brackish) waters. Liman Lagoon, which is connected to the Black Sea, has salinity levels between 0.24‰ and 13.19 ‰ (Macun, 2014). Salinity in Akyatan Lagoon in the northeastern Mediterranean region

varies between 3.3‰ and 35.34‰ (average 15.64 ± 6.4‰) (Akyol & Manaşırılı, 2023). According to Akyol and Ceyhan (2010), several carp specimens accidentally caught in 2004 were recorded from Peso Lagoon (Edirne, northern Aegean Sea), where salinity dropped to 0.04‰–0.09‰ in December during the rainy season. Among these lagoons, Homa is the most saline, with values ranging from 35.6‰ to 63.7‰ (Başdemir, 2017). Perdikaris *et al.* (2012) and references therein reported that *C. gibelio* was distributed in many continental freshwaters (lotic and lentic ecosystems) as well as transitional coastal waters with low salinity. Therefore, the occurrence of *C. gibelio* as a potamodromous fish in the highly saline Homa Lagoon represents an unexpected phenomenon. The area where the fish was observed may have lower salinity levels compared to other parts of the lagoon due to freshwater inputs through drainage canals.

C. gibelio individuals inhabiting lakes may migrate to river estuaries during winter to avoid low dissolved oxygen conditions (Froese & Pauly, 2025). Additionally, the heavy rainfall and flooding events in January and February 2026 may have facilitated the transport of this fish to the study area. It is therefore possible that the specimen reached the estuary of the Gediz River and subsequently moved through the canals into the lagoon. A similar situation was reported from the Akçapınar Stream delta, Gökova Bay (southeastern Aegean Sea) in 2020, where a specimen of *Capoeta aydinensis* (Cyprinidae) was spotted at a depth of 1.5 m, having been transported into the marine environment during floods caused by heavy rainfall (Akyol *et al.*, 2020). Another possible explanation is foraging activity in brackish lagoons. Bohlen (1999) noted that several freshwater fishes migrate to brackish waters for feeding and growth but return to lower-salinity waters for spawning.

In conclusion, a *C. gibelio* individual, potentially displaced during flood events caused by heavy rainfall or having entered the area during foraging activity, has been recorded for the first time from an unexpected saline habitat. Therefore, this paper not only reports a remarkable occurrence of this non-indigenous species in an unusual environment, but also contributes to the fish fauna records of Homa Lagoon, Izmir Bay. Further surveys are needed to determine whether this record represents a single sporadic occurrence or a more persistent presence.

NEPRIČAKOVANI POJAV SREBRNEGA KORESLJA *CARASSIUS GIBELIO* (CYPRINIDAE) V LAGUNI HOMA (IZMIRSKI ZALIV, EGEJSKO MORJE)

Okan AKYOL & Halil ŞEN
Ege University Faculty of Fisheries, 35440 Urla, Izmir, Türkiye
e-mail: okan.akyol@ege.edu.tr

POVZETEK

V laguni Homa na obali Izmirskega zaliva je bila 24. februarja 2026 ujeta samička srebrnega koreslja (*Carassius gibelio*). Riba je bila ujeta na globini enega metra s troslojno stojno mrežo, namenjeno lovu cipljev. Pojav je nenavaden, saj je laguna Homa izrazito slano okolje, kjer se vrednosti slanosti gibljejo med 35,6 ‰ in 63,7 ‰. Srebrni koreselj je namreč potamodromna ribja vrsta, ki je v teh vodah ne bi pričakovali. Ta članek ne prikazuje le izjemnega pojava te invazivne vrste v nenavadnem habitatu, ampak prinaša tudi dodaten prispevek k zapisom o favni rib v laguni Homa v Izmirskem zalivu.

Ključne besede: slana laguna, nenavaden habitat, meritve, Sredozemlje

REFERENCES

- Acarlı, D., A. Kara, B. Bayhan & T. Çoker (2009):** Catch composition and catch yield of species caught from Homa Lagoon (Izmir Bay, Aegean Sea). *Ege J. Fish. & Aquat. Sci.*, 26(1), 39–47.
- Afanasyev, S., O. Gupalo, O. Lietytska, Y. Kovalenko & D. Bănăduc (2025):** First record of *Carassius gibelio* (Bloch, 1782) in the upper reaches of the Tergi River (Greater Caucasus): distribution and morphological characteristics in new high-altitude conditions. *BiolInvasions Records*, 14(4), 907–927.
- Akyol, O. (2005):** The Last Lagoon of Izmir Bay: Homa. *Ekoloji Magazin*, pp.12-16.
- Akyol, O. & T. Ceyhan (2010):** Fisheries in Enez Lagoon (Edirne, Northern Aegean). *Ege J. Fish. & Aquat. Sci.*, 27(1), 31–34. (in Turkish)
- Akyol, O., V. Ünal & H.M. Sarı (2020):** The first report on the phenomenon of *Capoeta aydinensis* (Cyprinidae), occurring in Gökova Bay, Aegean Sea. *Ege J. Fish. & Aquat. Sci.*, 37(4), 423–425.
- Akyol, B. & M. Manaşırılı (2023):** Some properties reproductive of Prussian carp, *Carassius gibelio* Bloch, 1982, in Akyatan Lagoon, Adana, Türkiye. *Acta Biologica Turcica*, 36(3), J8, 1–8.
- Alpbaz, A. & T. Kınacıgil (1988):** A study on the fish abundance and fish fauna of Homa Lagoon in İzmir. *Ege J. Fish. & Aquat. Sci.*, 5(17–18), 31–56. (in Turkish)
- Baran, I. & T. Ongan (1988):** Limnological characteristics of Gala Lake, fishing problems and recommendations. Gala Gölü ve Sorunları Sempozyumu, Doğal Hayatı Koruma Derneği Bilimsel Yayınlar Serisi, İstanbul, pp. 46–54. (in Turkish)
- Başdemir, D. (2017):** Benthic and planktonic dinoflagellates of Homa Lagoon (Izmir Bay) and their relationships with environmental conditions. *Ege Univ. Fen Bil. Enst. Ph.D. Thesis*, 94 pp. (in Turkish)
- Bohlen, J. (1999):** Influence of salinity on the early development in the spined loach, *Cobitis taenia*. *J. Fish Biol.*, 55, 189–198.
- Ekmekçi, F.G., Ş.G. Kırankaya, L. Gençoğlu & B. Yoğurtçuoğlu (2013):** Present status of invasive fishes in inland waters of Turkey and assessment of the effects of invasion. *İstanbul Üniversitesi Su Ürünleri Dergisi*, 28, 105–140. (in Turkish)
- Froese, R. & D. Pauly (2025):** FishBase. World Wide Web electronic publication. www.fishbase.org, version (11/2025).
- Macun, S. (2014):** Age, Growth and Sex Ratio of *Cyprinus Carpio* (Linnaeus, 1758) in a lagoon lake, Lake Karaboğaz (Samsun, Turkey). *Hacettepe J. Biol. & Chem.*, 42 (3), 361–371.
- Perdikaris, C., A. Ergolavou, E. Gouva, C. Nathanailides, A. Chantzarpoulos & I. Paschos (2012):** *Carassius gibelio* in Greece: the dominant naturalised invader of freshwaters. *Rev. Fish. Biol. Fisheries*, 22, 17–27.
- Soto, I., P. Balzani, L. Carneiro, R.N., Cuthbert, R. Macêdo, A.S. Tarkan, D.A. Ahmed, A. Bang, K. Bace-la-Spychalska, S.A. Bailey, T. Baudry, L. Ballesteros-Mejia, A. Bortolus, E. Briski, J.R. Britton, M. Buřič, M. Camacho-Cervantes, C. Cano-Barbacil, D. Copilaş-Ciocianu, N.E. Coughlan, P. Courtois, Z. Csabai, T. Dalu, V. De Santis, J.W. E. Dickey, R.D. Dimarco, J. Falk-Andersson, R.D. Fernandez, M. Florencio, A.C.S. Franco, E. García-Berthou, D. Giannetto, M.M. Glavendekic, M. Grabowski, G. Heringer, I. Herrera, W. Huang, K.L. Kamelamela, N.I. Kirichenko, A. Kouba, M. Kourantidou, I. Kurtul, G. Laufer, B. Liptak, C. Liu, E. Lopez-Lopez, V. Lozano, S. Mammola, A. Marchini, V. Meshkova, M. Milardi, D.L. Musolin, M.A. Nuñez, F.J. Oficialdegui, J. Patoka, Z. Pattison, D. Pincheira-Donoso, M. Piria, A.F. Probert, J. J. Rasmussen, D. Renault, F. Ribeiro, G. Rillov, T.B. Robinson, A.E. Sanchez, E. Schwindt, J. South, P. Stoett, H. Verreycken, L. Vilizzi, Y-J. Wang, Y. Watari, P.M. Wehi, A. Weiperth, P. Wiberg-Larsen, S. Yapıcı, B. Yoğurtçuoğlu, R.D. Zenni, B.S. Galil, J.T.A. Dick, J.C. Russell, A. Ricciardi, D. Simberloff, C.J.A. Bradshaw & P.J. Haubrock (2024):** Taming the terminological tempest in invasion science. *Biol. Rev.*, 99(4), 1357–1390.
- Uysal, İ. & B. Boz (2018):** The most dangerous invasive alien species in Turkey and poisonous marine alien species in Turkey Report, October 2018. T.C. Tarım ve Orman Bakanlığı Doğa Koruma ve Milli Parklar Genel Müdürlüğü, Ankara. 68 pp.
- Yerli, S.V., F. Mangıt, Ö. Emiroğlu, V. Yeğen, R. Uysal, E. Ünlü, A. Alp, E. Buhan, T. Yıldırım & M. Zengin (2014):** Distribution of invasive *Carassius gibelio* (Bloch, 1782) (Teleostei: Cyprinidae) in Turkey. *Turk. J. Fish. Aquat. Sci.*, 14, 581–590.