


ANNALES



*Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 28, 2018, 2*



UDK 5

ISSN 1408-533X (Print)
ISSN 2591-1783 (Online)



ANNALES

**Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies**

Series Historia Naturalis, 28, 2018, 2

KOPER 2018

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Lektor/Supervisione/Language editor:

Polona Šergon (sl.), Petra Berlot (angl.)

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Prelom/Composizione/Typesetting:

Grafis trade d.o.o.

Tisk/Stampa/Print:

Grafis trade d.o.o.

Izdajatelja/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 14. 12. 2018.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS), Luka Koper in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation: 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL).

Vsi članki so v barvni verziji prosti dostopni na spletni strani: <http://zdjp.si/p/annalesshn/>
All articles are freely available in color via website: <http://zdjp.si/en/p/annalesshn/>

VSEBINA / INDICE GENERALE / CONTENTS

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
MEDITERRANEAN SHARKS**Hakan KABASAKAL, Erdi BAYRI & EYLÜL ATAÇ**

- Recent records of the great white shark,
Carcharodon carcharias (Linnaeus, 1758)
(Chondrichthyes: Lamnidae), in Turkish
waters (eastern Mediterranean) 93
Recentni podatki o belem morsklem volku,
Carcharodon carcharias (Linnaeus, 1758)
(Chondrichthyes: Lamnidae), v turških vodah
(vzhodno Sredozemlje)

**Christian CAPAPÉ, Sihem RAFRAFI-NOUIRA,
Khadija OUNIFI-BEN AMOR & Mohamed
Mourad BEN AMOR**

- Additional records of sandbar shark,
Carcharhinus plumbeus (Chondrichthyes:
Carcharhinidae) from the northern Tunisian
coast (central Mediterranean Sea) 99
Novi zapisi o pojavljanju sivega morskega psa,
Carcharhinus plumbeus (Chondrichthyes:
Carcharhinidae) na severni tunizijski obali
(osrednje Sredozemsko morje)

RECENTNE SPREMENBE V SREDOZEMSKI
BIODIVERZITETI
CAMBIAMENTI RECENTI NELLA
BIODIVERSITÀ MEDITERRANEA
RECENT CHANGES IN MEDITERRANEAN
BIODIVERSITY**Christian CAPAPÉ, Jeanne ZAOUALI,
Khadija OUNIFI-BEN AMOR & Mohamed
Mourad BEN AMOR**

- First record of Red sea goatfish *Parupeneus
forsskali* (Osteichthyes: Mullidae) from
Tunisian waters (central Mediterranean sea) 107
Prvi zapis o pojavljanju bradača vrste
Parupeneus forsskali (Osteichthyes: Mullidae)
iz tunizijskih voda (osrednje Sredozemsko morje)

Thodoros E. KAMPOURIS & Ioannis E. BATJAKAS

- The northernmost record of the thermophilic
Mediterranean parrotfish *Spurisoma cretense*
(Linnaeus, 1758) (Perciformes, Scaridae)
in the eastern Mediterranean Sea
(northwestern Aegean Sea) 111
Najsevernejši zapis o pojavljanju topoljubne
morske papige *Spurisoma cretense*
(Linnaeus, 1758) (Perciformes, Scaridae)
v vzhodnem Sredozemskem morju
(severo Zahodno Egejsko morje)

IHTIOLOGIJA
ITTOLOGIA
ICHTHYOLOGY**İlker AYDIN & Okan AKYOL**

- Occurrence of pearl fish, *Carapus acus*
(Osteichthyes: Carapidae) in Çeşme, Izmir
(Aegean Sea, Turkey) 119
Pojavljanje strmorinca, *Carapus acus*
(Osteichthyes: Carapidae) v predelu Çeşme,
Izmir (Egejsko morje, Turčija)

**Mohamed Mourad BEN AMOR, Khadija
OUNIFI-BEN AMOR & Christian CAPAPÉ**

- Additional records and extension of the
range of blackfish, *Centrolophus niger*
(Osteichthyes: Centrolophidae) from the
Tunisian coast (central Mediterranean Sea) 123
Novi zapis o pojavljanju črnuha, *Centrolophus
niger* (Osteichthyes: Centrolophidae)
iz tunizijske obale (osrednje Sredozemsko morje)

Lovrenc LIPEJ, Domen TRKOV & Borut MAVRIČ

- Occurrence of ribbon fish (*Trachipterus
trachypterus*) in Slovenian waters
(northern Adriatic Sea) 129
Pojavljanje kosice (*Trachipterus trachypterus*)
v slovenskem morju (severni Jadran)

FLORA
FLORA
FLORA**Nina REUTOVA, Petimat DZHAMBETOVA & Serekbay ABILEV**

- Species of wild flora as indicators of environmental genotoxicity 137
Rastlinske vrste kot indikatorji okoljske genotoksičnosti

Amelio PEZZETTA

- Le Orchidaceae del Comune di Pisino (Pazin, Croazia) 147
Kukavičevke pazinske občine (Pazin, Hrvaška)

FAVNA
FAUNA
FAUNA**Ana FORTIČ & Borut MAVRIČ**

- First record of the bryozoan *Tricellaria inopinata* (d'Hondt and Occhipinti-Ambrogi, 1985) from the Slovenian sea 155
Prvi zapis o pojavljanju mahovnjaka Tricellaria inopinata (d'Hondt and Occhipinti-Ambrogi, 1985) iz slovenskega morja

Olga Valentinovna GRISHAEVA & Kulyash Baizukevna KALIEVA

- Macrozoobenthos of arid watercourses of Kazakhstan: the Ilek River case 161
Makrozoobentos aridnih vodnih teles v Kazahstanu: primer reke Ilek

MISCELLANEA

Matjaž URŠIČ, Matija KRIŽNAR & Pavel JAMNIK

- Pregled pleistocenske favne in analiza ugrizov na kosteh v jamah Bele vode nad Gorenjo Trebušo in Smoganicu nad Mostom na Soči 173
Review of Pleistocene fauna and the analysis of bone bite marks in the caves Bele vode near Gorenja Trebuša and Smogаница near Most na Soči

OCENE IN POROČILA
RECENSIONI E RELAZIONI
REVIEWS AND REPORTS**Lovrenc LIPEJ & Martina ORLANDO-BONACA**

- Book review: La biologia marina a Trieste e nell'Alto Adriatico 193

IN MEMORIAM

- In memory of Mark Hines (1950-2018)
(Jadran Faganeli) 201
 Navodila avtorjem 203
Istruzioni per gli autori 205
Instruction to Authors 207

- Kazalo k slikam na ovitku 210
Index to images on the cover 210

short scientific article
received: 2018-08-27

DOI 10.19233/ASHN.2018.13

THE NORTHERNMOST RECORD OF THE THERMOPHILIC MEDITERRANEAN PARROTFISH *SPARISOMA CRETENSE* (LINNAEUS, 1758) (PERCIFORMES, SCARIDAE) IN THE EASTERN MEDITERRANEAN SEA (NORTHWESTERN AEGEAN SEA)

Thodoros E. KAMPOURIS & Ioannis E. BATJAKAS

Marine Sciences Department, School of the Environment, University of the Aegean, University Hill, Mytilene,
Lesvos Island, 81100, Greece
E-mail: mard16012@marine.aegean.gr

ABSTRACT

On July 2018 a specimen of the thermophilic fish Sparisoma cretense, commonly known as Mediterranean parrotfish, was captured in the Thermaic Gulf in the northwestern Aegean Sea, Greece, constituting the first such record in the region. The species mainly inhabits the southeastern shores of the Mediterranean, and its presence is in line with previous records of other thermophilic species in the northern Aegean which can be associated with the rise of sea temperatures as one of the signs of climate change. This record is the northernmost in the Eastern Mediterranean and the second northernmost record for the whole Mediterranean basin.

Key words: *Sparisoma cretense*, thermophilic species, new records, biodiversity, climate change, Aegean Sea

RITROVAMENTO PIÙ A NORD DEL PESCE PAPPAGALLO MEDITERRANEO TERMOFILO *SPARISOMA CRETENSE* (LINNAEUS, 1758) (PERCIFORMES, SCARIDAE) NEL MEDITERRANEO ORIENTALE (MAR EGEO NORD-OCCIDENTALE)

SINTESI

Nel luglio 2018 un esemplare del pesce termofilo Sparisoma cretense, comunemente noto come pesce pappagallo mediterraneo, è stato catturato nel Golfo Termico nel mar Egeo nord-occidentale, in Grecia, costituendo il primo ritrovamento della specie nella regione. La specie vive principalmente lungo le coste sud-orientali del Mediterraneo e la sua presenza è in linea con le precedenti segnalazioni di altre specie termofile nell'Egeo settentrionale, che possono venir associate all'innalzamento della temperatura del mare come uno dei segni del cambiamento climatico. Questo ritrovamento è il più settentrionale del Mediterraneo orientale e il secondo più settentrionale per l'intero bacino del Mediterraneo.

Parole chiave: *Sparisoma cretense*, specie termofile, nuovi ritrovamenti, biodiversità, cambiamenti climatici, mar Egeo

INTRODUCTION

The Mediterranean parrotfish, *Sparisoma cretense* (Linnaeus, 1758) is the only native parrotfish species of the Mediterranean Sea and can be added to the records of a few other temperate species within the Scaridae family (Froese & Pauly, 2018). So far, two alien Scaridae species have been reported within the Mediterranean, *Chlorurus rhakoura* Randall & Anderson, 1997 off the Italian coast (Insacco & Zava, 2017), and *Scarus ghobban* Forsskål, 1775 reported for the first time off the coasts of Israel in 2002 (Goren & Aronov, 2002).

The Mediterranean parrotfish occurs along the east coast of the Atlantic Ocean from Senegal to Portugal and within the Mediterranean Sea, mainly inhabiting the south and east coasts (Lousky, 2015). Similar to *Thalassoma pavo* (Linnaeus, 1758), the Mediterranean parrot fish is an ancient Mediterranean colonizer of Atlantic origin (Domingues et al., 2008). It can be found at a maximum depth of 50 m (Petrakis & Papaconstantinou, 1990), but more usually between 5–15 m of depth (Guidetti & Boero, 2002). Mainly active during the day (Azzurro et al., 2007; 2013), it is a herbivorous species feeding on algae and seagrasses, e.g. *Posidonia oceanica* (Linnaeus Delile, 1813 (de Girolamo et al., 1999) and *Cymodocea nodosa* (Ucria) Ascherson, 1870 (Del Río et al., 2016). While the adult population prefers mixed habitats with seagrass meadows and reefs, juveniles display no such preferences (Espino et al., 2015). The species commonly measures from 14 to 32 cm (total length) (Petrakis & Papaconstantinou, 1990), but might grow as big as 52 cm (total length) (Filiz & Sevingel, 2015).

The Mediterranean parrotfish has a moderate longevity and is a slow-growing fish that may reach up to 8 years of age (Petrakis & Papaconstantinou, 1990). It spawns from July to September (Petrakis & Papaconstantinou, 1990; Guidetti & Boero, 2002), displays sexual dichromatism (de Girolamo et al., 1999) and a dual mating system, with either multi-male groups or one dominant male holding harems (Alfonso et al., 2002). The species has low economic importance but is locally exploited in the Dodecanese (Petrakis & Papaconstantinou, 1990).



Fig. 1: The male *S. cretense* specimen captured off the Thermaic Gulf.
Sl. 1: Samec morske papige ujet v zalivu Thermaikos.

nou, 1990). Recent records from the northern Mediterranean coast, off Italy (Azzurro et al., 2011), from the Adriatic Sea, Croatia (Kruschel et al., 2012), from the northwest Greek Ionian Sea (Perdikaris et al., 2012), from Provence, France (Astruch et al., 2016), and from the Genoa Gulf, Italy (Bianchi et al., 2017), as well as observed expansions in the Aegean Sea, on both Greek and Turkish coasts (Papaconstantinou, 2014; Yapıcı et al., 2016), also suggest a correlation with the sea temperature rise and climate change (Fogarty et al., 2017).

MATERIAL AND METHODS

A single adult male *S. cretense* (Fig. 1) was caught on 23 July 2018 in the Thermaic Gulf, at approximately 40° 04' 10.7" N and 23° 19' 59.3" E (Fig. 2), by trammel nets with a mesh size of 22 mm, on hard bottom mixed with *P. oceanica* meadows. The depth ranged from 30 to 35 m. Morphometric and meristic data were obtained. Lengths (Tab. 1) were measured with the use of digital callipers to the nearest 0.01 mm. Weight (Tab. 1) was measured with an electronic scale to the nearest gram. The specimen was deposited at the Ichthyology laboratory of the Department of Marine Sciences, University of the Aegean.

RESULTS AND DISCUSSION

The identification was based on the criteria description by Golani et al. (2006) and Lousky (2015). Table 1 presents the related morphometric and meristic data. This study reports the northernmost record in the eastern Mediterranean Sea as acknowledged by F.A.O (2018), and the second northernmost record in the whole basin



Fig. 2: Map of the Thermaic Gulf, NW Aegean Sea, indicating sampling location (black dot).
Sl. 2: Zemljevid obravnavanega območja (zaliv Thermaikos, SZ Egejsko morje) z označeno lokaliteto ulova morske papige (črna pika).

(see Kruschel *et al.*, 2012 for details). The species was described by the fishermen as “never before seen,” which further supports the importance of this finding.

The Aegean Sea has a complex topography (Olson *et al.*, 2007). The area displays important physiochemical variations between north and south (Zervakis & Georgopoulos, 2002). The northern Aegean Sea is an area of high productivity compared to the eastern Mediterranean, due to its interconnection with the Black Sea (Petihakis *et al.*, 2014). Also, the northern Aegean Sea exhibits unique bottom topography features (the North Aegean Trough) that divide the area into three sub-basins: the Lemnos-Saros Islands Basin in the east, the Athos Basin in the centre, and the North Sporades Basin in the west (Zervakis & Georgopoulos, 2002, Olson *et al.*, 2007).

Recently some thermophilic fish species, native and alien, were reported off the northern Aegean Sea, with Kampouris *et al.* (2013) reporting the first record of oilfish, *Ruvettus pretiosus* Cocco, 1833, from the Chalkidiki Peninsula. Minos & Economidis (2015) report further occurrences of the tripletail *Lobotes surinamensis* (Bloch, 1790) from the Thermaic Gulf. The most striking is perhaps the record of the alien fish *Plectrohinichus gaterinus* (Forsskål, 1775) from the Thracian Sea, with no further records along the southeast Aegean and Levantine coasts (Corsini-Foka & Sarlis, 2016), which presents similarities with the first report of the African surgeonfish *Acanthurus monroviae* Steindachner, 1876 in Hellenic waters (Batjakas *et al.*, 2015).

CONCLUSIONS

The in-depth knowledge and experience of fishers has been recognized as a valuable source of scientific information that should be further acknowledged and utilized. Wider collaboration should be established among scientists, practitioners and policy makers (Azzuro *et al.*, 2018). Moreover, the contribution of naturalists and citizen-scientists on biodiversity monitoring

Tab. 1: Morphometric and meristic characters of the *S. cretense* specimen captured off the Thermaic Gulf.

Tab. 1: Morfometrija in meristični znaki pri morski papigi, ujeti v zalivu Thermaikos.

Morphometric characters	Value
Total length	262.57 mm
Standard length	218.78 mm
Head length	65.36 mm
Snout length	24.27 mm
Eye diameter	11.76 mm
Body depth	71.14 mm
Depth of caudal peduncle	34.24 mm
Max body width	34.94 mm
Min body width	26.57 mm
Weight	317 g
Meristic counts	
Dorsal fin	XI+10
Anal fin	III+9
Pectoral fin	12
Pelvic fin	I+5

(e.g., Kampouris *et al.*, 2018a) and on the detection and expansion of invasive alien species is already highly acknowledged (e.g. Kondylatos *et al.*, 2017; Giovos *et al.*, 2018; Kampouris *et al.*, 2018b) and wider stakeholder synergies should be established.

ACKNOWLEDGEMENTS

Both authors are grateful to professional fishermen Dimitrios and Kyriakos Ganigiannis, for providing the specimen in view and sharing further information.

NAJSEVERNEJŠI ZAPIS O POJAVLJANJU TOPLOLJUBNE MORSKE PAPIGE *SPARISOMA CRETENSE* (LINNAEUS, 1758) (PERCIFORMES, SCARIDAE) V VZHODNEM SREDOZEMSKEM MORJU (SEVEROZAHODNO EGEJSKO MORJE)

Thodoros E. KAMPOURIS & Ioannis E. BATJAKAS

Marine Sciences Department, School of the Environment, University of the Aegean, University Hill, Mytilene, Lesvos Island, 81100, Greece
E-mail: mard16012@marine.aegean.gr

POVZETEK

Julija 2018 je bil v zalivu Thermaikos v severozahodnem Egejskem morju (Grčija) ujet primerek morske papige Sparisoma cretense, kar predstavlja prvi primer pojavljanja te vrste v regiji. Vrsta domuje pretežno v predelih jugovzhodnih obal Sredozemlja. Njeno pojavljanje je tako kot v primerih pojava drugih topoljubnih vrst v severnem Egejskem morju, povezano z dvigom temperature morja in podnebnimi spremembami. Gre za najsevernejši podatek o pojavljanju te vrste v vzhodnem Sredozemlju in drugi najsevernejši podatek za celoten sredozemski bazen.

Ključne besede: *Sparisoma cretense*, termofilna vrsta, nov zapis, biodiverziteta, podnebne spremembe, Egejsko morje

REFERENCES

- Alfonso, P., T. Morato, & R.S. Santos (2002):** Spatial patterns in reproductive traits of the temperate parrotfish *Sparisoma cretense*. Fish Res., 90, 92–99. DOI: 10.1016/j.fishres.2007.09.029
- Astruch, P., P. Bonhomme, A. Goujard, E. Rouanet, C. Boudouresque, J. Harmelin & M. Harmelin-Vivien (2016):** Provence and Mediterranean warming: The parrotfish *Sparisoma cretense* is coming. Rapp. Comm. int. Mer Médit., 41, 362.
- Azzurro, E., A. Pais, P. Consoli & F. Andaloro (2007):** Evaluating day–night changes in shallow Mediterranean rocky reef fish assemblages by visual census. Mar Biol. 151, 2245–2253. DOI:10.1007/s00227-007-0661-9
- Azzurro, E., G. Agzzi, F. Maynou, J.J. Chiesa, & D. Savini (2013):** Diel rhythms in shallow Mediterranean rocky-reef fishes: a chronobiological approach with the help of trained volunteers. J. Mar. Biol. Assoc. U.K., 93(2), 461–470. DOI: 10.1017/S0025315412001166
- Azzurro, E., L. Bolognini, B. Dragičević, D. Drakulović, J. Dulčić, E. Fanelli, F. Grati, J. Kolitari, L. Lipej, E. Magaletti, O. Marković, S. Matic-Skoko, B. Mavrič, N. Milone, A. Joksimović, J. Tomanić, A. Scarpati, P. Tutman, D., Vrdoljak & F. Zappacosta (2018):** Detecting the occurrence of indigenous and non-indigenous megafauna through fishermen knowledge: A complementary tool to coastal and port surveys. Mar. Pollut. Bull., In Press. DOI: 10.1016/j.marpolbul.2018.01.016
- Azzurro, E., P. Moschella & F. Maynou (2011):** Tracking Signals of Change in Mediterranean Fish Diversity Based on Local Ecological Knowledge. PLoS ONE, 6(9), e24885. DOI: 10.1371/journal.pone.0024885
- Batjakas, I.E., T.E. Kampouris & A. Papadas (2015):** First Record of the African Surgeonfish *Acanthurus monroviae* (Osteichthyes: Acanthuridae) In the Hellenic Waters. JAMB, 2(6), 00047. DOI: 10.15406/jamb.2015.02.00047
- Bianchi, C.N., F. Caroli, P. Guidetti & C. Morri (2017):** Seawater warming at the northern reach for southern species: Gulf of Genoa, NW Mediterranean. J. Mar. Biol. Assoc. U.K., 98(1), 1–12. DOI: 10.1017/S0025315417000819
- Corsini-Foka, M. & N. Sarlis (2016):** A strange occurrence of *Plectrohinchus gaterinus* (Actinopterygii: Perciformes: Haemulidae) in the Thracian Sea (Eastern Mediterranean). Acta Ichthyol. Piscat., 46(1), 37–41. DOI: 10.3750/AIP2016.46.1.05
- de Girolamo, M., M. Scaggiante & M.B. Rasotto (1999):** Social organization and sexual pattern in the Mediterranean parrotfish *Sparisoma cretense* (Teleostei: Scaridae). Mar. Biol., 135(2), 353–360.
- Del Río, L., J. Vidal, S. Betancor & F. Tuya (2016):** Differences in herbivory intensity between the seagrass *Cymodocea nodosa* and the green alga *Caulerpa prolifera* inhabiting the same habitat. Aquat. Bot., 128, 48–57. DOI: 10.1016/j.aquabot.2015.10.001
- Domingues, V.S., M. Alexandrou, V.C. Almada, D.R. Robertson, A. Brito, R.S. Santos & G. Bernardi (2008):** Tropical fishes in a temperate sea: evolution of the wrasse *Thalassoma pavo* and the parrotfish *Sparisoma cretense* in the Mediterranean and the adjacent Macaronesian and Cape Verde Archipelagos. Mar. Biol., 154(3), 465–474.
- Espino, F., J.A. González, R. Haroun & F. Tuya (2015):** Abundance and biomass of the parrotfish *Sparisoma cretense* in seagrass meadows: temporal and spatial differences between seagrass interiors and seagrass adjacent to reefs. Environ. Biol. Fish., 98, 121–133. DOI 10.1007/s10641-014-0241-z
- F.A.O (2018):** Mediterranean and Black Sea (Major Fishing Area 37). World Wide Web electronic publication, www.fao.org/fishery/area/Area37/en (11/2018)
- Filiz, H. & N. Sevingel (2015):** A new maximum length for the parrotfish *Sparisoma cretense* (Linnaeus, 1758) in the Mediterranean Sea. J. Aquacult. Eng. Fish Res., 1(3), 140–143. DOI: 10.3153/JAEFR15015
- Fogarty, H.E., Burrows, M.T., Pecl, G.T., Robinson, L.M. & E.S. Poloczanska (2017):** Are fish outside their usual ranges early indicators of climate-driven range shifts? Glob. Change Biol., 23, 2047–2057. DOI: 10.1111/gcb.13635
- Froese, R. & D. Pauly (2018):** FishBase. World Wide Web electronic publication, www.fishbase.org, version (8/2018).
- Giovos, I., B. Giacomo, G. Romanidis-Kyriakidis, D. Marmara & P. Kleitou (2018):** First records of the fish *Abudedefduf sexfasciatus* (Lacepède, 1801) and *Acanthurus sohal* (Forsskål, 1775) in the Mediterranean Sea. Bioinvasions Rec., 7(2), 205–210. DOI: 10.3391/bir.2018.7.2.14
- Golani, D., B. Öztürk & N. Basusta, (2006):** Fishes of the Eastern Mediterranean. Turkish Marine Research Foundation, Istanbul, 260 pp.
- Goren, M. & A. Aronov (2002):** First record of the Indo-Pacific Parrot fish *Scarus ghobban* in the eastern Mediterranean. Cybium, 26, 239–240.
- Guidetti, P & F. Boero (2002):** Spatio-temporal variability in abundance of the parrotfish, *Sparisoma cretense*, in SE Apulia (SE Italy, Mediterranean Sea). Ital. J. Zool., 69 (3), 229–232. DOI: 10.1080/11250000209356464
- Insacco, G. & B. Zava (2017):** *Chlorurus rhakoura* Randall & Anderson, 1997 (Perciformes, Scaridae), an Indo-Pacific fish new to the Mediterranean Sea. Med. Mar. Sci., 18(2), 285–291.
- Kampouris, T.E., F. Tiralongo, A. Golemaj, I. Giovos, N. Doumpas & I.E. Batjakas (2018b):** *Penaeus aztecus* Ives, 1891 (Decapoda, Dendrobranchiata, Penaeidae): On the range expansion in Sicilian waters and on the first record from Albanian coast. Int. J. Fish. Aquat. Stud., 6(4), 468–471.
- Kampouris, T.E., I.E. Batjakas & I. Nikolopoulou (2013):** Occurrence of oilfish *Ruvettus pretiosus* Cocco, 1829 (Perciformes: Gempylidae) in Chalkidiki peninsula, North Aegean Sea, Greece. J. Biol. Res-Thessalon., 19, 165–167

Kampouris, T.E., V. Kouloumperis & I.e. Batjaksas (2018a): First documented record of the Palaemonid shrimp *Brachycarpus biunguiculatus* (Lucas, 1846) (Decapoda, Caridea, Palaemonidae) from Aegean Sea, Greece. Spixiana. In Press.

Kondylatos G., T.E. Kampouris, V. Kouloumperis & M. Corsini-Foka (2017): The Indo-Pacific brachyuran *Charybdis (Gonioinfradens) paucidentatus* (A. Milne-Edwards, 1861) (Brachyura, Portunidae) in the Cyclades, Aegean Sea. Tyrk. J. Zool., 41(6), 1118-1120.

Kruschel, C., I. Zubak & S. Schultz (2012): New records of the parrot fish, *Sparisoma cretense* and the cleaver wrasse, *Xyrichtys novacula*, by visual census in the southern Adriatic. Annales, Ser. Hist. Nat. 22(1), 47-53.

Louisy, P. (2015): Europe and Mediterranean marine fish. Identification Guide. Ulmer, Paris. pp. 182-183.

Minos, G. & P.S. Economidis (2015): More records on the tripletail, *Lobotes surinamensis* (Bloch, 1790) from the Aegean Sea. Front. Mar. Sci., 00056

Olson, D.B., V.H. Kourafalou, W.E. Johns, G. Samuels & M. Veneziani (2007): Aegean surface circulation from a satellite -tracked drifter array. J. Phys. Oceanogr., 37(7), 1898-1917.

Papaconstantinou, C. (2014): Fauna Graeciae. An updated checklist of the fishes in the Hellenic Seas, Monographs on Marine Sciences, 7, HCMR, Athens, 340 pp.

Perdikaris, C., E. Konstantinidis & I. Paschos (2012): *Sparisoma cretense* (L. 1758) from the NW Ionian coast of Greece. p. 164. In: Nicolaïdou et al., 2012. New Mediterranean Marine biodiversity records (June 2012). Mediterr. Mar. Sci., 13(1), 162-174.

Petihakis G., K. Tsiaras, G. Triantafyllou, S. Kalaroni & A. Pollani (2014): Sensitivity of the N. Aegean Sea ecosystem to the Black Sea Water inputs. Mediterr. Mar. Sci., 15(4), 790-804.

Petrakis, G. & C. Papaconstantinou (1990): Biology of *Sparisoma cretense* in the Dodecanese (Greece). J. Appl. Ichthyol., 6, 14-23.

Yapıcı, S., H. Filiz & G. Bilge (2016): Northwards range expansion of *Sparisoma cretense* (Linnaeus, 1758) in the Turkish Aegean Sea. J. Aquacult. Eng. Fish Res., 2(4), 201-207.

Zervakis, V. & D. Georgopoulos (2002): Hydrology and circulation in the North Aegean (eastern Mediterranean) throughout 1997 and 1998. Mediterr. Mar. Sci., 3(1), 5-19.