

ANNALES



*Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 29, 2019, 2*



UDK 5

ISSN 1408-533X (Print)
ISSN 2591-1783 (Online)



ANNALES

**Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies**

Series Historia Naturalis, 29, 2019, 2

KOPER 2019

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitzsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Lektor/Supervisione/Language editor:

Polona Šergon (sl.), Petra Berlot Kužner (angl.)

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelja/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria[®]

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente[®]

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 21. 12. 2019.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS), Luka Koper in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation: 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL).

Vsi članki so v barvni verziji prosti dostopni na spletni strani: <http://zdjp.si/p/annalesshn/>
All articles are freely available in color via website: <http://zdjp.si/en/p/annalesshn/>

VSEBINA / INDICE GENERALE / CONTENTS 2019(2)

50 LET MORSKE BIOLOŠKE POSTAJE
 NACIONALNEGA INŠITUTA ZA BIOLOGIJO
I 50 ANNI DELLA STAZIONE DI BIOLOGIA
MARINA DELL'ISTITUTO NAZIONALE
DI BIOLOGIA
50 YEARS OF THE MARINE BIOLOGY STATION
OF THE NATIONAL INSTITUTE OF BIOLOGY

Lovrenc LIPEJ, Domen TRKOV, Domen STANIČ,**Sara CERNICH & Saul CIRIACO**

First Record of Sergeant Major, *Abudefduf saxatilis* (Linnaeus, 1758) in the Adriatic Sea 159
Prvi zapis o pojavljanju velikega seržanta, Abudefduf saxatilis (Linnaeus, 1758)
 v Jadranskem morju

Valentina PITACCO & Selahattin Ünsal KARHAN

First Record of *Flabelliderma cinari* Karhan,
 Simboura & Salazar-Vallejo, 2012 (Polychaeta:
Flabelligeridae) from the Adriatic Sea 167
Prvi zapis o pojavljanju vrste Flabelliderma
cinari Karhan, Simboura & Salazar-Vallejo, 2012
 (*Polychaeta: Flabelligeridae*) v Jadranskem morju

**Ana FORTIČ, Domen TRKOV, Borut MAVRIČ
& Lovrenc LIPEJ**

Assessment of Bryozoan Xenodiversity
 in the Slovenian Coastal Sea 173
Ocena ksenodiverzitete mahovnjakov
 v slovenskem obalnem morju

**Martina ORLANDO-BONACA, Janja FRANCÉ,
Borut MAVRIČ & Lovrenc LIPEJ**

Impact of the Port of Koper on
Cymodocea nodosa Meadow 187
Vpliv koprskega pristanišča na travnik
kolenčaste cimodoceje (*Cymodocea nodosa*)

IGHTIOLOGIJA
ITTOLOGIA
ICHTHYOLOGY

**Akraam Faraj KARA, Mohamed AL HAJAJI,
Hisham GHMATI & Esmail A. SHAKMAN**

Food and Feeding Habits of *Mustelus mustelus*
 (Linnaeus, 1758) (Chondrichthyes: Triakidae)
 Along the Western Coast of Libya 197
Prehranjevalne navade navadnega morskega
psa, Mustelus mustelus (Linnaeus, 1758)
 (Chondrichthyes: Triakidae), vzdolž zahodne
 obale Libije

**Youssouph DIATTA, Almamy DIABY, Sihem
RAFRAFI-NOUIRA & Christian CAPAPÉ**

Capture of a Rare Endangered Species Leafscale
Gulper Shark *Centrophorus squamosus*
 (Chondrichthyes: Squalidae) from the Coast
 of Senegal (Eastern Tropical Atlantic) 205
Ulov redkega in ogroženega morskega psa
Centrophorus squamosus (Chondrichthyes:
Squalidae) iz obalnih voda Senegala (vzhodni
 tropski Atlantik)

**Sihem RAFRAFI-NOUIRA, Christian REYNAUD
& Christian CAPAPÉ**

Morphological Deformities in a Striped Sea
 Bream *Lithognathus mormyrus* (Osteichthyes:
Sparidae) from Northern Tunisian Waters
 (Central Mediterranean Sea) 211
Morfološke deformacije pri ovčici Lithognathus
mormyrus (Osteichthyes: Sparidae) iz severno
 tunizijskih vod (osrednje Sredozemske morje)

Sihem RAFRAFI-NOUIRA, Christian REYNAUD & Christian CAPAPÉ

- On a Occurrence of *Gadella maraldi*
(Osteichthyes: Gadiformes: Moridae) on the
Tunisian Coast (Central Mediterranean Sea) 219
O pojavljanju vrste Gadella maraldi
(Osteichthyes: Gadiformes: Moridae) ob
tunizijski obali (osrednje Sredozemsko morje)

Mohamed Mourad BEN AMOR, Marouène BDIOUI, Khadija OUNIFI-BEN AMOR, Christian REYNAUD & Christian CAPAPÉ

- Unusual Record of Round Fantail Stingray
Taeniurus grabata (Chondrichthyes:
Dasyatidae) from the Tunisian Coast
(Central Mediterranean Sea) 223
Nenavadni zapis o pojavljanju morskega biča
vrste *Taeniurus grabata* (Chondrichthyes:
Dasyatidae) iz tunizijskih voda (osrednje
Sredozemsko morje)

**JADRANSKA MORSKA FAVNA
FAUNA MARINA ADRIATICA
ADRIATIC MARINE FAUNA**

- Roland R. MELZER, Martin HEß, Manuel A. STAGGL, Tihomir MAKOVEC & Borut MAVRIČ**
Hippolyte prideauxiana Leach, 1817: First
Record for the Northern Adriatic and
Observations on Mimetic Colouration 231
Hippolyte prideauxiana Leach, 1817: prvi
zapis o pojavljanju v severnem Jadranu
s podatki o barvnemu prikrivanju

- Mona REZAEI, Al VREZEC, Borut MAVRIČ & Lovrenc LIPEJ**
Biometry and Population Gender Structure
of Three Crab Species (Crustacea:
Decapoda) from Sandy Bottom
in the Northern Adriatic Sea 235
Biometrija in spolna struktura pri treh vrstah
rakovic (Crustacea: Decapoda) iz sedimentnega
 dna severnega Jadranu

**ONESNAŽEVANJE OKOLJA
INQUINAMENTO DELL'AMBIENTE
ENVIRONMENTAL POLLUTION**

- Sezginer TUNCER, Sedat GÜNDÖĞDU, Cem ÇEVIK & Aytuğ ZILIFLI**
Belone belone (Linnaeus, 1760) and *Spicara smaris* (Linnaeus, 1758) Entangled in Plastic
Collars in the Dardanelles Strait, Turkey 247
Iglica, Belone belone (Linnaeus, 1760),
in girica, Spicara smaris (Linnaeus, 1758),
zapleteni v plastični ovoj v ožini
Dardanele, Turčija

- Jamila FLIOU, Ali AMECHROUQ, Mohammed ELHOURRI, Ouassima RIFFI & Mostafa EL IDRISI**
Determination of Heavy Metal Content
in a *Daphne gnidium* L. Plant Using Atomic
Absorption Spectroscopy 253
Določevanje vsebnosti težkih kovin v volčinu
vrste *Daphne gnidium* L. z uporabo atomske
absorpcijske spektroskopije

**DELO NAŠIH ZAVODOV IN DRUŠTEV
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS AND ASSOCIATIONS**

- Simon KERMA**
Half a Century of the Marine
Biology Station Piran 261
Pol stoletja delovanja Morske
biološke postaje Piran

- Navodila avtorjem 267
Istruzioni per gli autori 269
Instruction to Authors 271
Kazalo k slikam na ovitku 273
Index to images on the cover 273

received: 2019-10-17

DOI 10.19233/ASHN.2019.20

CAPTURE OF A RARE ENDANGERED SPECIES LEAFSCALE GULPER SHARK *CENTROPHORUS SQUAMOSUS* (CHONDRICHTHYES: SQUALIDAE) FROM THE COAST OF SENEGAL (EASTERN TROPICAL ATLANTIC)

Youssouph DIATTA & Almamy DIABY

Laboratoire de Biologie marine, Institut fondamental d'Afrique noire, (IFAN Ch. A. Diop), Université Cheikh Anta Diop de Dakar, BP 206, Dakar, Senegal

Sihem RAFRAFI-NOUIRA

Unité de Recherches Exploitation des Milieux aquatiques, Institut Supérieur de Pêche et d'Aquaculture de Bizerte, Université de Carthage, BP 15, 7080 Menzel Jemil, Tunisia

Christian CAPAPÉ

Laboratoire d'Icthyologie, case 104, Université de Montpellier, 34 095 Montpellier cedex 5, France
 e-mail: capape@univ-montp2.fr

ABSTRACT

*This note presents a recent record of a specimen of rare shark leafscale gulper shark *Centrophorus squamosus*. A description of the specimen is given, including morphometric measurements, total body weight and dental formulae. Comments on this unusual and rare capture off the Senegalese coast are also provided.*

Keywords: Squalidae, distribution, conditions of capture, dental formula, coast of Senegal

CATTURA DI UNA SPECIE RARA E MINACCIATA, IL CENTROFORO SQUAME A FOGLIA *CENTROPHORUS SQUAMOSUS* (CHONDRICHTHYES: SQUALIDAE), AL LARGO DELLA COSTA DEL SENEGAL (ATLANTICO TROPICALE ORIENTALE)

SINTESI

*Questa nota tratta una cattura recente di un esemplare di una rara specie di squali, il centroforo squame a foglia, *Centrophorus squamosus*. Viene fornita una descrizione dell'esemplare, comprese le misurazioni morfometriche, il peso corporeo totale e le formule dentali. Gli autori inoltre discutono questa insolita e rara cattura al largo della costa senegalese.*

Parole chiave: Squalidae, distribuzione, condizioni di cattura, formula dentale, costa senegalese

INTRODUCTION

The leafscale gulper shark *Centrophorus squamosus* (Bonnaterre, 1788) is widely distributed on both sides of the Indian Ocean and off the western Pacific coast (White, 2003). The species is known along the eastern Atlantic shore from Iceland and the Atlantic Slope to the Canary Islands, Senegal, the Faeroes, Madeira and the Azores (White, 2003). Southward, *C. squamosus* occurs from Senegal, Gulf of Guinea to the west coast of South Africa (McEachran & Branstetter, 1984).

Previously, a total of 37 specimens *Centrophorus squamosus* had been examined, among them two large pregnant females carrying near term embryos (Cadenat & Blache, 1981). Since then, no specimen had been recorded although investigations were continuously and regularly conducted in the area (Capapé et al., 1994,

2001; Diatta et al., 2009; Diatta, 2014). Such lack of captures suggested that the species probably no longer occurred or, at least, had disappeared from the coast of Senegal. Conversely, in the wake of collaboration with experienced local fishermen, we were informed that a specimen of *Centrophorus squamosus* had been captured and delivered to our laboratory for a thorough examination. This new record forms the subject of the present paper, which includes a description of the specimen and some comments about the rare occurrence of this species in Senegalese waters.

MATERIAL AND METHODS

On 19 February 2015, a specimen of *Centrophorus squamosus* was caught by means of gill nets on rocky bottoms at a depth of 150–250 m, off the fishing site of Ouakam, located 5 km north of Dakar, in the Cape Verde Peninsula ($17^{\circ}36'04.20''W$ and $14^{\circ}40'39.31''N$) (Fig. 1).

The specimen was measured to the nearest millimetre and weighed to the nearest gram. Morphological measurements and teeth counts on upper and lower jaws were carried out following Compagno (1984) and are summarized in Table 1. The specimen was fixed in buffered formaldehyde and deposited in the Ichthyological Collection of the Institute Cheikh Anta Diop of Dakar (Senegal), under catalogue number IFAN Centr-squa 01.

RESULTS AND DISCUSSION

The studied specimen was a female, measuring 1300 mm in total length and weighing 11,200 g in total body weight (TWB). It was identified as a *Centrophorus squamosus* based on a combination of main morphological characters, such as: snout slightly long, broadly parabolic; no anal fin; both dorsal fins with large spines; first dorsal lower than the second dorsal, the latter moderately large, as high as or higher than first dorsal fin; free rear tips of pectoral fins, margins slightly extended (Fig. 2A); upper teeth erect at the centre third, but not symmetrical; lower teeth with serrated edges (Fig. 2B); denticles on the sides of the body closely-set and overlapping, leaf-like, with a high median ridge and a small lateral ridge on each side (Fig. 2C); colour uniformly dark grey.

The general morphology, the shapes of teeth and denticles, the morphometric measurements and the dental formula are in total agreement with Bass et al. (1976), Cadenat & Blache (1981), Compagno (1984) and McEachran & Branstetter (1984), confirming the identification of this shark, which constitutes the latest known record of *C. squamosus* from the coast of Senegal.

Species of the genus *Centrophorus* Müller & Henle, 1817 are generally targeted by fishermen, especially for the extraction of liver oil, which is highly appreciated in cosmetology and pharmacology (Diatta, 2014). Nowadays, only specimens of gulper shark *C. granulosus* (Bloch & Schneider, 1801) are commonly captured




Fig. 1: Map of the coast of Senegal indicating (black star) the capture site of the specimen of *Centrophorus squamosus* (ref. IFAN Centr-squa 01), redrawn from Diatta et al. (2013).

Sl. 1: Zemljevid senegalske obale z označbo lokalitete, kjer je bil primerek (ref. IFAN Centr-squa 01) vrste *Centrophorus squamosus* ujet. Pritejeno po Diatta s sod. (2013).

Tab 1: Morphometric measurements (in mm and as % TL), total body weight (in gram) and dental formulae recorded in the specimen of *Centrophorus squamosus* (ref. IFAN Centr-squa 01).**Tab. 1: Morfometrične meritve (v mm in kot % dolžine telesa), celotna telesna teža (v gramih) in zobna formula primerka vrste *Centrophorus squamosus* (ref. IFAN Centr-squa 01).**

Reference	IFAN Centr-squa 01	
Sex	Female	
Total body weight (g)	11200	
Measurements	mm	% TL
Total length	1300	100.0
Fork length	1140	87.7
Precaudal length	1030	79.2
Pre-first dorsal length	43	3.3
Pre-second dorsal length	830	63.8
Head length	290	22.3
Prebranchial length	220	16.9
Respiracular length	150	11.5
Preorbital length	60	4.6
Prepectoral length	253	19.5
Prepelvic length	780	60.0
Snout-vent length	880	67.7
Interdorsal space	310	23.8
Dorsal caudal space	81	6.2
Pectoral-pelvic space	500	38.5
Pelvic-caudal space	700	53.8
Vent-caudal length	460	35.4
Eye length	50	3.8
Eye height	30	2.3
Prenarial length	45	3.5
Intergill length	60	4.6
First gill slit height	35	2.7
Fifth gill slit height	42	3.2
Pectoral anterior margin	141	10.8
Pectoral inner margin	88	6.8
Pectoral posterior margin	103	7.9
Pectoral height	152	11.7
Dorsal caudal margin	210	16.2
Terminal caudal lobe	122	9.4
Subterminal caudal margin	27	2.1
Preventral caudal margin	168	12.9
First dorsal anterior margin	190	14.6

First dorsal height	80	6.2
First dorsal posterior margin	135	10.4
First dorsal base	205	15.8
First dorsal length	280	21.5
First dorsal inner margin	90	6.9
Second dorsal anterior margin	130	10.0
Second dorsal height	42	3.2
Second dorsal posterior margin	103	7.9
Second dorsal inner margin	35	2.7
Second dorsal base	125	9.6
Head height	180	13.8
Trunk height	210	16.2
Abdomen height	190	14.6
Tail height	110	8.5
Caudal peduncle height	55	4.2
Pelvic midpoint-second dorsal insertion	230	17.7
First dorsal midpoint- pelvic origin	270	20.8
Second dorsal insertion- pelvic insertion	140	10.8
Pelvic posterior margin length	80	6.2
Pelvic inner margin length	65	5.0
Pelvic base	87	6.7
Pelvic length	145	11.2
Pelvic anterior margin	130	10.0
Pelvic height	103	7.9
Interorbital space	102	7.8
Head width	240	18.5
Trunk width	300	23.1
Abdomen width	350	26.9
Caudal peduncle width	30	2.3
Mouth length	50	3.8
Mouth width	190	14.6
Internarial space	40	3.1
Dental formulae	16-1-16/13-1-13	




Fig. 2: A. General morphology of the specimen of *Centrophorus squamosus* (ref. IFAN Centr-squa 01) caught off the coast of Senegal, scale bar = 200 mm. B. Jaw of the same specimen showing upper and lower teeth, scale bar = 100 mm. C. Denticles removed from the dorsal surface of the same specimen, scale bar = 2 mm.
Sl. 2: A. Morfologija primerka vrste *Centrophorus squamosus* (ref. IFAN Centr-squa 01), ujetega ob obali Senegala, merilo = 200 mm. B. Čeljust z zgornjimi in spodnjimi zobmi, merilo = 100 mm. C. Dentikli s hrbitne površine raziskanega primerka, merilo = 2 mm.

throughout the Senegalese coast (Diatta et al., 2009). Some morphological differences allow distinguishing *C. squamosus* from *C. granulosus*. *C. squamosus* displays free rear tips of the pectoral fins, broadly angular and not reaching past the first dorsal spine, denticles in adults exhibit multiple lateral cusps. In *C. granulosus*, the free rear tips of the pectoral fins extend into narrow angular lobes, which reach past the first dorsal fin origin, and denticles in adults are without cusps, broadly rounded posteriorly. Since local fishermen are familiar with captures of *C. granulosus* in the area, they were able to distinguish this species from *C. squamosus*, and therefore they could immediately bring it to the laboratory for a thorough examination. The rarity of the species in the area could also explain why these fishermen could express a positive, or at least neutral, attitude to this shark species. The specimen was captured over continental shelf grounds at a depth of 150 to 200 m, which is normally shallower than the depth ranges

reported by McEachran & Branstetter (1984). According to McEachran & Branstetter (1984), *C. squamosus* is a benthic species inhabiting deeper areas, from 400 to 1875 m, rarely below 1000 m. Although these areas are poorly exploited by fishermen, this does not fully explain the scarcity of the species in commercial catches. However, this isolated capture of a single specimen suggests that the species is not completely extinct in the area, but other records are needed to state if a viable population still occurs in the area or if the species is dangerously threatened.

ACKNOWLEDGMENTS

The authors wish to thank the fishermen from the fishing site of Ouakam, Senegal, who provided the specimen, and Dr Mafalda Freitas, Head of the Marine Biology Station of Funchal, Madeira Island who enabled us to identify the specimen as *Centrophorus squamosus*.

ULOV REDKEGA IN OGROŽENEGA MORSKEGA PSA *CENTROPHORUS SQUAMOSUS*
(CHONDRICHTHYES: SQUALIDAE) IZ OBALNIH VODA SENEGALA
(VZHODNI TROPSKI ATLANTIK)

Youssouph DIATTA & Almamy DIABY

Laboratoire de Biologie marine, Institut fondamental d'Afrique noire, (IFAN Ch. A. Diop), Université Cheikh Anta Diop de Dakar, BP 206, Dakar, Senegal

Sihem RAFRAFI-NOUIRA

Unité de Recherches Exploitation des Milieux aquatiques, Institut Supérieur de Pêche et d'Aquaculture de Bizerte, Université de Carthage, BP 15, 7080 Menzel Jemil, Tunisia

Christian CAPAPÉ

Laboratoire d'Ichtyologie, case 104, Université de Montpellier, 34 095 Montpellier cedex 5, France
e-mail: capape@univ-montp2.fr

POVZETEK

Avtorji poročajo o ulovu primerka redkega morskega psa vrste *Centrophorus squamosus*. Navajajo opis vrste, morfometrične meritve, telesno težo in zobno formulo. Poleg tega razpravljajo o tem redkem in nenavadnem ulov ob obali Senegala.

Ključne besede: Squalidae, razširjenost, podatki o ulovu, zobna formula, senegalska obala

REFERENCES

- Bass, A.J., J.D D'Aubrey & N. Kistnasamy (1976):** Sharks of the east coast of southern Africa. Part V: The Families Oxynotidae, Chlamydoselachidae, Squalidae, Dalatiidae and Echinorhinidae. Invest. Rep. Oceanogr. Res. Inst. Durban, 45, 1-103.
- Cadenat, J. & J. Blache (1981):** Requins de Méditerranée et d'Atlantique (plus particulièrement de la côte occidentale d'Afrique). Faune tropicale, ORSTOM, 21, 1-330.
- Capapé, C., M. Diop & M. N'Dao (1994):** Observations sur dix-sept espèces de Sélaïens d'intérêt économique capturées dans la région marine de Dakar-Ouakam (Sénégal, Atlantique tropical oriental). Bull. Inst. fond. Afr. noire Cheikh Anta Diop, Dakar, 1994, sér. A, 47, 87-102.
- Capapé, C., A. Gueye-Ndiaye, Y. Diatta, M. Diop & A.A. Seck (2001):** Observations on six elasmobranch species recorded from off the coast of Senegal (eastern tropical Atlantic). Acta Adriat., 42(1), 89-102.
- Compagno, L.J.V. (1984):** FAO Species Catalogue, vol. 4, Sharks of the World. An annotated and illustrated catalogue of shark species known to date. FAO Fisheries Synopsis, 125, vol. 4, part 1 (non carcharhinoids): viii+1–250 pp.
- Diatta, Y., A. Gueye-Ndiaye, P. Ndiaye, S. Ndaw, C. Reynaud & C. Capapé (2009):** Production of elasmobranch species off the coast of Senegal (eastern tropical Atlantic). Elasmovisor, 4, 22-25.
- Diatta, Y., C. Reynaud & C. Capapé (2013):** First case of albinism recorded in striped panray, *Zanobatus schoenleinii* (Chondrichthyes: Platyrrhinidae) from the coast of Senegal (Eastern Tropical Atlantic). J. Ichthyol., 53(11), 1007-1012.
- Diatta, Y. (2014):** Contribution to the knowledge of the bioecology of 26 elasmobranch species from the coast of Senegal Thesis, Dakar, University Cheikh Anta Diop of Dakar, Senegal, 402 pp.
- Mc Eachran, J.D. & S. Branstetter (1984):** Squalidae. In: P.J.P. Whitehead, M.L. Bauchot, J.C. Hureau., J. Nielsen J.& E. Tortonese. (Editors), pp. 128-147. Fishes of the North-western Atlantic and the Mediterranean, Vol I, UNESCO, Paris.
- White, W.T. (2003):** *Centrophorus squamosus*. The IUCN Red List of Threatened Species. <https://www.iucnredlist.org/species/41871/10581731>. Accessed on 16 October 2019.