

ANNALES



*Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 29, 2019, 2*



UDK 5

ISSN 1408-533X (Print)
ISSN 2591-1783 (Online)



ANNALES

**Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies**

Series Historia Naturalis, 29, 2019, 2

KOPER 2019

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitzsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Lektor/Supervisione/Language editor:

Polona Šergon (sl.), Petra Berlot Kužner (angl.)

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelja/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria[®]

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente[®]

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 21. 12. 2019.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS), Luka Koper in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation: 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL).

Vsi članki so v barvni verziji prosti dostopni na spletni strani: <http://zdjp.si/p/annalesshn/>
All articles are freely available in color via website: <http://zdjp.si/en/p/annalesshn/>

VSEBINA / INDICE GENERALE / CONTENTS 2019(2)

50 LET MORSKE BIOLOŠKE POSTAJE
 NACIONALNEGA INŠITUTA ZA BIOLOGIJO
I 50 ANNI DELLA STAZIONE DI BIOLOGIA
MARINA DELL'ISTITUTO NAZIONALE
DI BIOLOGIA
50 YEARS OF THE MARINE BIOLOGY STATION
OF THE NATIONAL INSTITUTE OF BIOLOGY

**Lovrenc LIPEJ, Domen TRKOV, Domen STANIČ,
 Sara CERNICH & Saul CIRIACO**
 First Record of Sergeant Major, *Abudefduf saxatilis* (Linnaeus, 1758) in the Adriatic Sea 159
Prvi zapis o pojavljanju velikega seržanta, Abudefduf saxatilis (Linnaeus, 1758)
v Jadranskem morju

Valentina PITACCO & Selahattin Ünsal KARHAN
 First Record of *Flabelliderma cinari* Karhan, Simboura & Salazar-Vallejo, 2012 (Polychaeta: Flabelligeridae) from the Adriatic Sea 167
Prvi zapis o pojavljanju vrste Flabelliderma cinari Karhan, Simboura & Salazar-Vallejo, 2012 (Polychaeta: Flabelligeridae) v Jadranskem morju

Ana FORTIČ, Domen TRKOV, Borut MAVRIČ & Lovrenc LIPEJ
 Assessment of Bryozoan Xenodiversity in the Slovenian Coastal Sea 173
Ocena ksenodiverzitete mahovnjakov v slovenskem obalnem morju

Martina ORLANDO-BONACA, Janja FRANCÉ, Borut MAVRIČ & Lovrenc LIPEJ
 Impact of the Port of Koper on *Cymodocea nodosa* Meadow 187
Vpliv koprskega pristanišča na travnik kolenčaste cimodoceje (*Cymodocea nodosa*)

IGHTIOLOGIJA
ITTOLOGIA
ICHTHYOLOGY

Akraam Faraj KARA, Mohamed AL HAJAJI, Hisham GHMATI & Esmail A. SHAKMAN
 Food and Feeding Habits of *Mustelus mustelus* (Linnaeus, 1758) (Chondrichthyes: Triakidae) Along the Western Coast of Libya 197
Prehranjevalne navade navadnega morskega psa, Mustelus mustelus (Linnaeus, 1758) (Chondrichthyes: Triakidae), vzdolž zahodne obale Libije

Youssouph DIATTA, Almamy DIABY, Sihem RAFRAFI-NOUIRA & Christian CAPAPÉ
 Capture of a Rare Endangered Species Leafscale Gulper Shark *Centrophorus squamosus* (Chondrichthyes: Squalidae) from the Coast of Senegal (Eastern Tropical Atlantic) 205
Ulov redkega in ogroženega morskega psa Centrophorus squamosus (Chondrichthyes: Squalidae) iz obalnih voda Senegala (vzhodni tropski Atlantik)

Sihem RAFRAFI-NOUIRA, Christian REYNAUD & Christian CAPAPÉ
 Morphological Deformities in a Striped Sea Bream *Lithognathus mormyrus* (Osteichthyes: Sparidae) from Northern Tunisian Waters (Central Mediterranean Sea) 211
Morfološke deformacije pri ovčici Lithognathus mormyrus (Osteichthyes: Sparidae) iz severno tunizijskih vod (osrednje Sredozemske morje)

Sihem RAFRAFI-NOUIRA, Christian REYNAUD & Christian CAPAPÉ

- On a Occurrence of *Gadella maraldi*
(Osteichthyes: Gadiformes: Moridae) on the
Tunisian Coast (Central Mediterranean Sea) 219
O pojavljanju vrste Gadella maraldi
(Osteichthyes: Gadiformes: Moridae) ob
tunizijski obali (osrednje Sredozemsko morje)

Mohamed Mourad BEN AMOR, Marouène BDIOUI, Khadija OUNIFI-BEN AMOR, Christian REYNAUD & Christian CAPAPÉ

- Unusual Record of Round Fantail Stingray
Taeniurus grabata (Chondrichthyes:
Dasyatidae) from the Tunisian Coast
(Central Mediterranean Sea) 223
Nenavadni zapis o pojavljanju morskega biča
vrste *Taeniurus grabata* (Chondrichthyes:
Dasyatidae) iz tunizijskih voda (osrednje
Sredozemsko morje)

**JADRANSKA MORSKA FAVNA
FAUNA MARINA ADRIATICA
ADRIATIC MARINE FAUNA**

- Roland R. MELZER, Martin HEß, Manuel A. STAGGL, Tihomir MAKOVEC & Borut MAVRIČ**
Hippolyte prideauxiana Leach, 1817: First
Record for the Northern Adriatic and
Observations on Mimetic Colouration 231
Hippolyte prideauxiana Leach, 1817: prvi
zapis o pojavljanju v severnem Jadranu
s podatki o barvnemu prikrivanju

- Mona REZAEI, Al VREZEC, Borut MAVRIČ & Lovrenc LIPEJ**
Biometry and Population Gender Structure
of Three Crab Species (Crustacea:
Decapoda) from Sandy Bottom
in the Northern Adriatic Sea 235
Biometrija in spolna struktura pri treh vrstah
rakovic (Crustacea: Decapoda) iz sedimentnega
 dna severnega Jadranu

**ONESNAŽEVANJE OKOLJA
INQUINAMENTO DELL'AMBIENTE
ENVIRONMENTAL POLLUTION**

- Sezginer TUNCER, Sedat GÜNDÖĞDU, Cem ÇEVIK & Aytuğ ZILIFLI**
Belone belone (Linnaeus, 1760) and *Spicara smaris* (Linnaeus, 1758) Entangled in Plastic
Collars in the Dardanelles Strait, Turkey 247
Iglica, Belone belone (Linnaeus, 1760),
in girica, Spicara smaris (Linnaeus, 1758),
zapleteni v plastični ovoj v ožini
Dardanele, Turčija

- Jamila FLIOU, Ali AMECHROUQ, Mohammed ELHOURRI, Ouassima RIFFI & Mostafa EL IDRISI**
Determination of Heavy Metal Content
in a *Daphne gnidium* L. Plant Using Atomic
Absorption Spectroscopy 253
Določevanje vsebnosti težkih kovin v volčinu
vrste *Daphne gnidium* L. z uporabo atomske
absorpcijske spektroskopije

**DELO NAŠIH ZAVODOV IN DRUŠTEV
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS AND ASSOCIATIONS**

- Simon KERMA**
Half a Century of the Marine
Biology Station Piran 261
Pol stoletja delovanja Morske
biološke postaje Piran

- Navodila avtorjem 267
Istruzioni per gli autori 269
Instruction to Authors 271
Kazalo k slikam na ovitku 273
Index to images on the cover 273

received: 2019-10-24

DOI 10.19233/ASHN.2019.23

UNUSUAL RECORD OF ROUND FANTAIL STINGRAY *TAENIUROPS GRABATA* (CHONDRICHTHYES: DASYATIDAE) FROM THE TUNISIAN COAST (CENTRAL MEDITERRANEAN SEA)

Mohamed Mourad BEN AMOR & Marouène BDIOUI

Institut National des Sciences et Technologies de la Mer, port de pêche, 2025 La Goulette, Tunisia

Khadija OUNIFI-BEN AMOR

Université de Tunis El Manar, Faculté des Sciences de Tunis, Laboratoire de Biodiversité, Biotechnologie et Changement Climatique, LR11ES09, 1002, Tunis, Tunisia

Christian REYNAUD & Christian CAPAPÉ

Laboratoire d'Ictyologie, Université de Montpellier, case 104, 34095 Montpellier cedex 5, France
e-mail: capape@univ-montp2.fr

ABSTRACT

*This paper reports the capture of a specimen of round fantail stingray *Taeniurops grabata* (Geoffroy Saint-Hilaire, 1817) on the northern coast of Tunisia, caught in deep waters, at a depth of 450 m. It was a large specimen of 1.12 m in disc width, 2.70 m in total length and 110 kg in total body weight. This is the largest and heaviest specimen known to date in Tunisian waters, and probably in the Mediterranean and even outside it. Additionally, comments are given about the distribution of the species, well-known in the eastern Mediterranean Basin and eastern tropical Atlantic, but unknown in the western Mediterranean Basin.*

Key words: Dasyatidae, distribution, expansion range, Mediterranean Sea, eastern tropical Atlantic

RITROVAMENTO INUSUALE DI TRIGONE AFRICANO *TAENIUROPS GRABATA* (CHONDRICHTHYES: DASYATIDAE) LUNGO LA COSTA DELLA TUNISIA (MEDITERRANEO CENTRALE)

SINTESI

*L'articolo riporta la cattura di un esemplare di trigone africano, *Taeniurops grabata* (Geoffroy Saint-Hilaire, 1817), lungo la costa settentrionale della Tunisia, catturato in acque profonde, ossia a 450 m di profondità. L'esemplare era molto grande, con 1,12 m di larghezza del disco, 2,70 m di lunghezza totale e 110 kg di peso corporeo totale. Si tratta dell'esemplare più grande e più pesante ritrovato fino ad oggi nelle acque tunisine, e probabilmente anche nel Mediterraneo e al di fuori di esso. Gli autori inoltre discutono la distribuzione della specie, ben nota nel bacino del Mediterraneo orientale e nell'Atlantico tropicale orientale, ma sconosciuta nel bacino del Mediterraneo occidentale.*

Parole chiave: Dasyatidae, distribuzione, area di espansione, mare Mediterraneo, Atlantico tropicale orientale

INTRODUCTION

Round fantail stingray *Taeniurops grabata* (Geoffroy Saint-Hilaire, 1817) is found off the west African coast from Mauritania to Angola, around the São Tiago and Cape Verde Islands (Capapé & Desoutter, 1990). It has also been reported from the Azores Islands, where large free-swimming specimens were observed in the wild in shallow coastal waters (Barreiros, *pers. comm.*, 2019). *T. grabata* occurs throughout southeastern Mediterranean (McEachran & Capapé, 1984), especially from Turkish waters (Bilecenoglu *et al.*, 2014) to the Levant Basin (Golani, 2005), with the Syrian coast constituting to date the easternmost extension of its range (Ali *et al.*, 2013).

Taeniurops grabata has been reported in southern Tunisia, mainly in the Gulf of Gabès where a viable population appears to be successfully established (Capapé, 1989; Bradai *et al.*, 2004). However, more recently, Boudaya *et al.* (2018) studied parasitic infestation occurring in specimens captured off Bizerte, a city located in northern Tunisia. Additionally, a large specimen of *T. grabata* was captured from the same area and landed at the fish market of Zarzouna-Bizerte, where it was observed and photographed. This specimen is described in the present paper and some comments are given about the species distribution in the local area, the Mediterranean Sea and in the eastern tropical Atlantic.

MATERIAL AND METHODS

On 28 September 2019, a female specimen of *Taeniurops grabata* was caught by biface bottom trawl in the coastal waters of northern Tunisia, off Bizerte at 37°




Fig. 1: Map of northern Tunisia with the black star indicating the capture site of *Taeniurops grabata*. GT = Gulf of Tunis; GH=Gulf of Hammamet.

Sl. 1: Zemljovid severne Tunizije s crno zvezdico, ki označuje lokalitetu ulova vrste *Taeniurops grabata*. GT = tuniški zaliv; GH=hamameški zaliv.

40' N and 9° 90' E (Fig. 1). The capture occurred at a depth of 450 m, on soft bottom, unfortunately no information was provided about the associated fauna. Some measurements were recorded to the nearest centimetre (cm) and total body weight to the nearest kilogram (kg). Other morphometric measurement could not be taken, because the specimen was cut into slices after landing to be sold at the fish market of Bizerte; its flesh is appreciated in the preparation and consumption of local dishes.

RESULTS AND DISCUSSION

The specimen was identified as *Taeniurops grabata* following Mc Eachran & Capapé (1984): disc rather circular, broader than long, tail short and compressed, with a membranous fold below; dorsal surface almost entirely rugose, with a row of spines from the middle of the disc to the spine; dorsal surface brown with dark blotches and irregular vermiculation, belly beige (Fig. 2).

McEachran & Capapé (1984) noted that the maximum disc width of *T. grabata* could reach 1 m, in total agreement with specimens recorded by Capapé (1989) from the Gulf of Gabès. The present specimen measured 1.12 m in disc width, 2.7 m in total length, and its total body weight reached 110 kg. This is therefore the largest and heaviest *Taeniurops grabata* recorded to date in Tunisian waters, probably throughout the Mediterranean and even outside this sea. Large elasmobranch species, generally sharks, occur off the northern coast of the Tunisian transitional area between the eastern and western basins (Soufi-Kechaou *et al.*, 2018; Capapé *et al.*, 2018), but this was the first time that as large a batoid species as *T. grabata* was found there, and furthermore, in deep waters. This species generally inhabits shallow coastal waters, not exceeding 50 m (Capapé, 1989). This capture of *T. grabata* at a depth of 450 m is rather unusual, probably occasional, but it could also explain the rarity of the species in deep areas that are poorly exploited by fisheries.

Taeniurops grabata used to be considered as a species having tropical affinities and only found in southern Tunisian regions (Postel, 1956). Conversely, Boudaya *et al.* (2018) studied parasitic infestation occurring in specimens of *T. grabata* captured northward from Bizerte. Such occurrence was probably due to migration toward northern regions. Similar patterns were reported for other elasmobranch species previously unknown in this latter area (El Kamel *et al.*, 2009; El Kamel-Moutalibi *et al.*, 2014; Rafrafi-Nouira *et al.*, 2015; Rafrafi-Nouira, 2016; Soufi-Kechaou *et al.*, 2018). These migrations are likely owed to the warming of the Mediterranean waters, including the Tunisian coast, due to global climate change (Francour *et al.*, 1994). Additionally, the present capture of a large specimen indicates that the occurrence of a northern population of *T. grabata* cannot be totally ruled out. It could be also considered as the species' northwesternmost extent of habitat range in the




Fig. 2: The *Taeniurops grabata* captured from the northern coast of Tunisia. Dorsal surface = A. posterior view, B. anterior view, C. anterior margin view. D. tail, black arrow indicating the place of the sting removed by fishermen.

Sl. 2: Primerek vrste *Taeniurops grabata*, ujet ob severni tunizijski obali. Zgornja površina - A. pogled od zadaj, B. pogled od spredaj, C. pogled sprednjega roba. D. rep, črna puščica označuje predel, s katerega so ribiči odstranili trn.

Mediterranean Sea, as the species does not occur in the western Basin (Mc Eachran & Capapé, 1984) and is not reported northward off the coast of France (Capapé et al., 2006). Additionally, among Maghreb shores, *T. grabata* is unknown off the Algerian coast (Refes et al., 2010) and off the Moroccan coast (Lloris & Rocabado, 1998).

Boudaya et al. (2018) found the same anthocephalid species in *T. grabata* from the Tunisian coast and in those from the Senegalese coast. Additionally, three species belonging to the monogenean genus *Heterocotyle* Scoot 1904 parasitized the gills of *T. grabata* from these two distinct areas (Neifar et al., 1999). Such discoveries

suggest that the distribution of *T. grabata* is not really fragmented and that the species' non-occurrence could be due to lack of sampling effort (Boudaya et al., 2018). *T. grabata* is known to inhabit shallow coastal waters (Capapé, 1989; Barreiros, pers. comm., 2019), however the capture of the studied specimen in deep waters confirms the opinion of Ajmenian & Powers (2014), who noted that batoid species are very mobile and prone to both large latitudinal and vertical migrations. Still, only molecular tools could indicate whether or not different populations of *T. grabata* occur in the Mediterranean Sea and in the eastern tropical Atlantic.

ACKNOWLEDGEMENTS

The authors wish to kindly thank Dr João Pedro Barreiros from the University of Azores, Portugal for information

about *Taeniurops grabata* from Azores Islands, and Prof Lassad Neifar and Dr Lobna Boudaya from the University of Sfax, Tunisia for sending an important paper.

NENAVADNI ZAPIS O POJAVLJANJU MORSKEGA BIČA VRSTE *TAENIUROPS GRABATA*
(CHONDRICHTHYES: DASYATIDAE) IZ TUNIZIJSKIH VODA
(OSREDNJE SREDOZEMSKO MORJE)

Mohamed Mourad BEN AMOR & Marouène BDIOUI

Institut National des Sciences et Technologies de la Mer, port de pêche, 2025 La Goulette, Tunisia

Khadija OUNIFI-BEN AMOR

Université de Tunis El Manar, Faculté des Sciences de Tunis, Laboratoire de Biodiversité, Biotechnologie et Changement Climatique,
LR11ES09, 1002, Tunis, Tunisia

Christian REYNAUD & Christian CAPAPÉ

Laboratoire d'Ictyologie, Université de Montpellier, case 104, 34095 Montpellier cedex 5, France
e-mail: capape@univ-montp2.fr

POVZETEK

Avtorji poročajo o ulovu morskega biča vrste *Taeniurus grabata* (Geoffroy Saint-Hilaire, 1817) iz globine 450 m ob severni tunizijski obali. Gre za večji primerek, ki je meril 1,12 m v premeru telesnega diska in tehtal 110 kg. To je doslej največji in najtežji zabeleženi primerek te vrste v tunizijskih vodah in verjetno tudi v sredozemskih vodah in širše. Avtorji še razpravljajo o razširjenosti vrste, ki je sicer dobro raziskana v vzhodnem Sredozemskem morju in tropskem delu vzhodnega Atlantika, dokaj neznana pa v zahodnem delu Sredozemlja.

Ključne besede: Dasyatidae, razširjenost, širjenje areala, Sredozemsko morje, vzhodni tropski Atlantik

REFERENCES

- Ajemian, M. & S. Powers (2014):** Towed-float satellite telemetry tracks large-scale movement and habitat connectivity and habitat of myliobatid stingrays. Environ. Biol. Fish., 97, 1067-1081.
- Ali, M., A. Saad, C. Reynaud & C. Capapé (2013):** First records of the Round Fantail Stingray, *Taeniura grabata* (Chondrichthyes: Dasyatidae), off the Syrian coast (eastern Mediterranean). Zool. Midd. East., 59(2), 176-178.
- Bilecenoglu, M., M. Kaya, B. Cihangir & E. Åiçek (2014):** An updated checklist of the marine fishes of Turkey. Turk. J. Zool., 38(6), 901-929.
- Boudaya, L., L. Neifar & L. Euzet (2018):** A new genus and three new species of Anthocephaliidae (Cestoda, Rhinebothriidae) from the round fantail stingray, *Taeniurus grabata* (Chondrichthyes, Dasyatidae) from the Mediterranean Sea and Atlantic Ocean. J. Helminthology, 1-11, <https://doi.org/10.1017/S002214X18001025>.
- Bradai, M.N., J.P Quignard, A. Bouain, O. Jarboui, A. Ouannes-Ghorbel, L. Ben Abdallah, J. Zaouali & S. Ben Salem (2004):** Ichtyofaune autochtone et exotique des côtes tunisiennes: Recensement et biogéographie. Cybium, 28(4), 315-328.
- Capapé, C. (1989):** Les Sélaciens des côtes méditerranéennes: aspects généraux de leur écologie et exemples de peuplements. Océanis, 15(3), 309-331.
- Capapé, C. & M. Desoutter (1990):** Dasyatidae. In: J.C. Quéro, J.C. Hureau, C. Karrer, A. Post & Saldanha L. (Editors), pp. 59-63. Check list of the fishes of the eastern tropical Atlantic, Vol. 1. Unesco 1, Paris.
- Capapé C., O. Guélorget, Y. Vergne & J.-P. Quignard (2006):** Skates and rays (Chondrichthyes) from waters off the Languedocian coast (southern France, northern Mediterranean). Annales Series Historia Naturalis, 16(2), 166-178.
- Capapé C., S. Rafrafi-Nouira, K. Ounifi-Ben Amor & M.M. Ben Amor (2018):** Additional records of sandbar shark, *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae) from the northern Tunisian coast (central Mediterranean Sea). Annales Series Historia Naturalis, 28(2), 99-104.
- El Kamel, O., N. Mnasri, J. Ben Souissi, M. Boumaïza, M. M. Ben Amor & C. Capapé (2009):** Inventory of elasmobranch species caught in the Lagoon of Bizerte (north-eastern Tunisia, central Mediterranean). Pan-Amer. J. Aquat. Sci., 4(4), 383-412.
- El Kamel-Moutalibi O., N. Mnasri-Sioudi., S. Rafrafi-Nouira., M. Boumaïza., C. Reynaud & C. Capapé (2014):** Additional records of a rare elasmobranch species, sharpnose seven-gill shark *Heptranchias perlo* (Hexanchidae) off the northern Tunisian coast (central Mediterranean). Annales Series Historia Naturalis, 24(2), 96-109.
- Francour, P., C. F. Boudouresque, J.G. Harmelin., M.L Harmelin-Vivien. & J.-P. Quignard (1994):** Are the Mediterranean waters becoming warmer? Mar. Poll. Bull., 28(9), 523-526.
- Golani, D. (2005):** Check-list of the Mediterranean Fishes of Israel. Zootaxa, 947, 1-200.
- Lloris, D. & J. Rocabado (1998):** Guide FAO d'identification des espèces pour les besoins de la pêche. Guide d'identification des ressources marines vivantes pour le Maroc. FAO, Rome, 263 pp.
- McEachran, J.D & C. Capapé (1984):** Dasyatidae. In: Whitehead, P.J.P., M.L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese (eds): Fishes of the North-eastern Atlantic and the Mediterranean, Vol. 1. Unesco, Paris, pp. 197-202.
- Neifar, L., L. Euzet & O.K. Ben Hassine (1999):** Trois nouveaux Heterocotyle (Monogena, Monocotylidae) parasites branchiaux de *Taeniura grabata* (Euselachii, Dasyatidae) en Tunisie. Zoosystema, 21, 157-170.
- Postel, E. (1956):** Les affinités tropicales de la faune ichthyologique du golfe de Gabès. Bull. Sta. Océanogr. Pêche Salammbô, 52, 47-58.
- Rafrafi-Nouira, S., O. El Kamel-Moutalibi, C. Reynaud, M. Boumaïza & C. Capapé (2015):** Additional and unusual captures of elasmobranch species from the northern coast of Tunisia (central Mediterranean). J. Ichthyol., 55(6), 337-345.
- Rafrafi-Nouira, S. (2016):** Catalogue raisonné des espèces de poissons capturées devant Ras Jebel et autres régions marines de Tunisie septentrionale: aspects morphologiques, biométriques et bio-écologiques. Thesis, University of Bizerte (Tunisia), 509 pp.
- Refes, W., N. Semahi, M. Boulahdid & J.-P. Quignard (2010):** Inventaire de la faune ichtyologique de la côte orientale algérienne (El Kala; Annaba; Skikda; Jilel; Bejaïa). Rapp. Comm. int. Mer Médit., 39, 646.
- Soufi-Kechaou, E., K. Ounifi Ben Amor, J. Ben Souissi, M.M. Ben Amor & C. Capapé (2018):** The capture of a large predatory shark, *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae), off the Tunisian coast (Central Mediterranean). Annales Series Historia Naturalis, 28(1), 23-287.