


ANNALES



*Analí za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 30, 2020, 2*



UDK 5

ISSN 1408-533X
e-ISSN 2591-1783



ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 30, 2020, 2

KOPER 2020

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitzsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

**Urednica/Redattrice/Editor:
Lektor/Supervisione/Language editor:**

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Petra Berlot Kužner (angl.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Martina Orlando-Bonaca (sl./it.)

Tisk/Stampa/Print:

Dušan Podgornik, Lovrenc Lipej

Založništvo PADRE d.o.o.

Izdajatelja/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria[®]

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente[®]

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;

e-mail: annales@mbss.org, internet: www.zdjp.si

Redakcija te številke je bila zaključena 21. 12. 2020.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS), Luka Koper in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation: 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: Clarivate Analytics: Biological Abstracts / BIOSIS Previews / Zoological Record; Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>. The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>. Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>.



VSEBINA / INDICE GENERALE / CONTENTS 2020(2)

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
MEDITERRANEAN SHARKS**Paraskevi K. KARACHLE, Caterina STAMOULI & Aikaterini DOGRAMMATI**

Review of the Sharpnose Sevengill Shark
Heptranchias perlo (Chondrichthyes:
Hexanchidae) in the Mediterranean:
Historical and Recent Data 131
Pregled o pojavljanju morskega psa
sedmoroškrgarja Heptranchias perlo
(Chondrichthyes: Hexanchidae) v Sredozemlju:
zgodovinski in recentni podatki

Hakan KABASAKAL

A Historical Catch of White Shark,
Carcharodon carcharias (Lamniformes:
Lamnidae), in the Sea of Marmara (Turkey)
from the 1950s 147
Zgodovinski pregled ulova belega morskega
volka, Carcharodon carcharias (Lamniformes:
Lamnidae), v Marmarskem morju (Turčija) iz
petdesetih let

Nexhip HYSOLAKOJ, Rigers BAKIU & Alen SOLDO

The First Record of Kitefin Shark
Dalatias licha in Albanian Waters 151
Prvi zapis o pojavljanju klinoplavutega
morskega psa Dalatias licha v albanskih vodah

Deniz ERGÜDEN, Mahmut İĞDE, Cemal TURAN, Deniz AYAS & Hakan KABASAKAL

Occurrence of a Large Bigeye Thresher Shark,
Alopias superciliosus (Lamniformes: Alopiidae),
in the Northeastern Levantine Sea (İskenderun
Bay, Eastern Mediterranean Sea, Turkey) 157
Pojavljanje velike velikooke morske lisice,
Alopias superciliosus (Lamniformes: Alopiidae),
v severovzhodnem Levantskem morju (zaliv
İskenderun, vzhodni Mediteran, Turčija)

Lovrenc LIPEJ, Janja FRANCÉ, Domen TRKOV, Borut MAVRIČ & Aleš BOLJE

The Occurrence and Status of Thresher Shark
(Alopias vulpinus) in Waters Off Slovenia 165
Pojavljanje in status navadnih morskih lisic
(*Alopias vulpinus*) v vodah Slovenije

Hakan KABASAKAL & Erdi BAYRI

First Record of a Young-Of-The-Year
Carcharodon carcharias in the
Strait of the Dardanelles 175
Prvi zapis o pojavljanju prvoletnega
belega morskega volka (*Carcharodon*
carcharias) v dardanski ozini

Francesco TIRALONGO, Clara MONACO & Alessandro DE MADDALENA

Report on a Great White Shark *Carcharodon*
carcharias Observed Off Lampedusa, Italy 181
Zapis o pojavljanju belega morskega volka
Carcharodon carcharias pri Lampedusi, Italija

Hakan KABASAKAL

A Leucistic White Shark, *Carcharodon*
carcharias (Lamniformes: Lamnidae),
from the Northern Aegean Sea, Turkey 187
Belični primerek belega morskega volka,
Carcharodon carcharias (Lamniformes:
Lamnidae), iz severnega Egejskega
morja, Turčija

IHTIOLOGIJA

ITTOLOGIA
ICHTHYOLOGY**Christian CAPAPÉ, Youssouph DIATTA, Almamy DIABY, Sihem RAFRAFI-NOUIRA & Christian REYNAUD**

Leucistic Piebald Striped Panray,
Zanobatus schoenleinii (Chondrichthyes:
Zanobatidae), from the Coast of Senegal
(Eastern Tropical Atlantic) 193
Belični primerek vrste Zanobatus
schoenleinii (Chondrichthyes:
Zanobatidae) iz Senegalske obale
(vzhodni tropski Atlantik)

Fatiha DALOUCHÉ, Lotfi BENSAHLA-TALET, Sidi Mohamed El Amine ABI AYAD & Ahmed BENSAHLA-TALET

Gonadal Development of Bogues,
Boops boops (Linnaeus, 1758),
from Oran Bay (Teleostei, Sparidae) 201
Razvoj gonad pri bukvi, *Boops boops*
(Linnaeus, 1758), iz Oranskega zaliva
(Teleostei, Sparidae)

Francesco TIRALONGO & Roberto PILLON	FLORA
New Distributional Records of <i>Gobius buccichi</i> (Pisces, Gobiidae) from the Mediterranean Sea and In Situ Comparisons with <i>Gobius incognitus</i> 215	FLORA
<i>Novi podatki o razširjenosti vrste glavača Gobius buccichi (Pisces, Gobiidae) iz Sredozemskega morja in in situ primerjave z vrsto Gobius incognitus</i>	FLORA
BIOINVAZIJA	Amelio PEZZETTA
BIOINVASIONE	Le Papaveraceae Juss. della Flora Italiana: Distribuzione Regionale e Considerazioni Fitogeografiche 245
BIOINVASION	<i>Družina Papaveraceae Juss. v italijanski flori: regionalna razširjenost in fitogeografska opredelitev</i>
İnci TÜNEY-KIZILKAYA & Okan AKYOL	Toufik CHEDADI, Omar IDRISI, Anas ELKHABLI, Youssef KHACHTIB, Abdelmajid HADDIOUI & Mohammed EL HANSALI
Occurrence of <i>Scarus ghobban</i> (Scaridae) at the Border of the Aegean Sea (Kaş, Turkey) 223	Morphological Characterization of <i>Orobanche crenata</i> in Carrots and Legumes (Faba Bean and Chickpea): Indications of Potential Genetic Differentiation Towards Host Plants 265
<i>Pojavljanje modroprogaste morske papige (Scaridae) na meji Egejskega morja (Kaş, Turkey)</i>	<i>Morfološka opredelitev vrste Orobanche crenata pri korenju in stročnicah (bob in čičerika): pokazatelji možne genetske diferenciacije proti gostiteljskim rastlinam</i>
Argyro ZENETOS & Anastasia MILIOU	OCENE IN POROČILA
Abudefduf cf. <i>saxatilis</i> in the Saronikos Gulf, Greece: Unaided Introduction or Human Aided Transfer? 227	RECENSIONI E RELAZIONI
Abudefduf cf. <i>saxatilis</i> v zalivu Saronikos (Grčija): nenameren ali nameren vnos?	REVIEWS AND REPORTS
JADRANSKA MORSKA BIODIVERZITETA	Lovrenc Lipej
BIODIVERSITÀ MARINA DELL'ADRIATICO	Book review: ICHTHYOFAUNA OF THE ADRIATIC SEA 283
ADRIATIC MARINE BIODIVERSITY	Lovrenc Lipej
Martina ORLANDO-BONACA & Domen TRKOV	Book review: KNIGHTS, BALLERINAS AND INVISIBLES: THE DECAPOD CRUSTACEANS OF THE BRIJUNI MARINE PROTECTED AREA 284
After More Than Forty-Five Years a New Finding of <i>Cystoseira foeniculacea</i> f. <i>latiramosa</i> in the Coastal Sea of Slovenia 233	Alenka Malej
<i>Nova najdba vrste Cystoseira foeniculacea f. latiramosa v obalnem morju Slovenije po več kot štiridesetih letih</i>	Book review: BIODIVERZITETA BIOGENIH FORMACIJ. ZAKLADNICA NARAVE SLOVENSKEGA MORJA 285
Valentina PITACCO	POPRAVKI IN DOPOLNILA
First Record of the Echiurid <i>Maxmuelleria gigas</i> (M. Müller, 1852) in Slovenian Waters (Northern Adriatic) 239	ERRATA CORRIGE
<i>Prvi zapis o pojavljanju zvezdaša Maxmuelleria gigas (M. Müller, 1852) v slovenskih vodah (severni Jadran)</i>	ERRATA & CORRIGENDA
	Corrigendum to “Capture of a Large Great White Shark, <i>Carcharodon carcharias</i> (Lamnidae) from the Tunisian Coast (Central Mediterranean Sea): a Historical and Ichthyological Event”, Annales, Series Historia Naturalis, 2020, 30(1), 9-14 289
	Kazalo k slikam na ovitku 290
	<i>Index to images on the cover</i> 290

received: 2020-09-09

DOI 10.19233/ASHN.2020.23

**LEUCISTIC PIEBALD STRIPED PANRAY, *ZANOBATUS SCHOENLEINII*
(CHONDRICHTHYES: ZANOBATIDAE), FROM THE COAST OF SENEGAL
(EASTERN TROPICAL ATLANTIC)**

Christian CAPAPÉ

Laboratoire d'Ichtyologie, Université de Montpellier, case 104, 34095 Montpellier cedex 5, France
e-mail: capape@univ-montp2.fr

Youssouph DIATTA & Almamy DIABY

Laboratoire de Biologie marine, Institut fondamental d'Afrique noire, (IFAN Ch. A. Diop), Université Cheikh Anta Diop de Dakar,
BP 206, Dakar, Senegal

Sihem RAFRAFI-NOUIRA

Université de Carthage, Unité de Recherches Exploitation des Milieux aquatiques, Institut Supérieur de Pêche et d'Aquaculture de Bizerte,
BP 15, 7080 Menzel Jemil, Tunisia

Christian REYNAUD

Laboratoire Interdisciplinaire en Didactique, Education et Formation, Université de Montpellier, 2 place Marcel Godechot, B.P. 4152,
34092 Montpellier cedex 5, France

ABSTRACT

*In the present paper, a case of leucism/piebaldism is reported in a striped panray *Zanobatus schoenleinii* (Müller & Henle, 1841) captured at a low depth off the coast of Senegal (eastern tropical Atlantic). This is the second case recorded to date for this species. The specimen was sub-adult male with developed but flexible claspers, which measured 495 mm total length and 279 mm disc width, and weighed 751 g. Comments on albinism in fishes are provided in this paper, especially those reported in batoid species.*

Key words: morphology, colour, pigmentation, morphometric measurements, batoid species

**ESEMPLARE LEUCISTICO DI *ZANOBATUS SCHOENLEINII* (CHONDRICHTHYES:
ZANOBATIDAE) AL LARGO DELLA COSTA DEL SENEGAL
(ATLANTICO TROPICALE ORIENTALE)**

SINTESI

*L'articolo riporta un caso di leucismo/piebaldismo in un esemplare di *Zanobatus schoenleinii* (Müller & Henle, 1841), catturato in acque poco profonde al largo della costa del Senegal (Atlantico tropicale orientale). Si tratta del secondo caso di questo tipo registrato fino ad oggi per tale specie. L'esemplare era un maschio sub-adulto, con pterigopodi sviluppati ma flessibili, di lunghezza totale pari a 495 mm, larghezza del disco di 279 mm e 751 g di peso. Gli autori discutono i casi di albinismo nei pesci, in particolare quelli riportati per le specie batoidi.*

Parole chiave: morfologia, colore, pigmentazione, misure morfometriche, specie batoidi

INTRODUCTION

The striped panray, *Zanobatus schoenleinii* (Müller & Henle, 1841), is an endemic species known from the eastern tropical Atlantic, where it only occurs in the area between southern Morocco (Lloris & Rocabado 1998), Mauritania (Maurin & Bonnet, 1970), Senegal (Diatta et al., 2013), Guinea-Bissau

(Sanches, 1991), and the Gulf of Guinea (Blache et al., 1970).

Zanobatus schoenleinii is both commonly and abundantly captured in handicraft fishery throughout the coast of Senegal (Cadenat, 1951; Séret & Opic, 1990; Capapé et al., 1995). The species inhabits shallow coastal waters not exceeding 50 m of depth (Capapé et al., 1995). In the wake of a




Fig. 1: Map of the Senegalese coast indicating the capture site of the leucistic specimen of *Zanobatus schoenleinii* (black star).

Sl. 1: Zemljevid senegalske obale z označeno lokaliteto, kjer je bil ujet belični primerek vrste *Zanobatus schoenleinii* (črna zvezdica).

collaboration with experienced fishermen, several *Z. schoenleinii* were collected in shallow coastal waters surrounding the touristic area of Dakar. All collected specimens were delivered to the laboratory for thorough examination, including the abnormal specimens described and commented in the present paper.

MATERIAL AND METHODS

A total of 105 specimens were captured on 25 May 2019 using trammel nets at a depth between 5 and 10 m on soft bottom, off Ouakam, a handicraft fishery site located 5 km north Dakar, in Cape Verde Peninsula, Senegal, at 14° 43' 26" N and 17° 29' 21" W (Fig. 1). Of these specimens, 104 were normal and one specimen abnormal, displaying non-pigmented areas on the dorsal surface.

All specimens were measured to the nearest millimetre and weighed to the nearest gram. The normal specimens ranged between 253 mm and 580 mm in total length (TL), 154 and 340 mm in disc width (DW), and weighed between 104 and 1433 g in total body weight (TBW). However, a sub-sample including 51 normal specimens and the single abnormal specimen was used in order to obtain a more visible view, allowing to clearly distinguish this abnormal specimen from the normal specimens. The subgroup ranged between 250 and 300 mm DW and weighed between 561 and 1346 g TBW.

Morphometric measurements recorded on one of the normal specimens and the abnormal specimens are presented in Table 1, following Diatta et al. (2013). Both specimens were fixed in 10% buffered formaldehyde, preserved in 75 % ethanol and deposited in the Ichthyological Collection of the Institut Supérieur d'Aquaculture et de Pêche of Bizerte (Tunisia), receiving catalogue numbers, ISPAB-Zan-sch-01 for the normal specimen and ISPAB-Zan-sch-02 for the abnormal specimen.

The relation between DW and TBW was used as a complement following Froese et al. (2011), including all specimens, normal and abnormal, to show if this latter would have been able to develop in the wild like normal specimens. This relation, $TBW = aDW^b$, was converted into its linear regression, expressed in decimal logarithmic coordinates, and

correlations were assessed by least-squares regression as: $\log TBW = \log a + b \log DW$. Significance of constant b differences was assessed according to the hypothesis of isometric growth if $b = 3$, positive allometry if $b > 3$, negative isometry if $b < 3$ (Pauly, 1983). These two latter tests were performed by using logistic model STAT VIEW 5.0.

RESULTS AND DISCUSSION

All normal specimens were identified as *Zanobatus schoenleinii* following the combination of main



Fig. 2: Normal specimen of *Zanobatus schoenleinii* (Ref. ISPAB- Zan-sch-01), from the coast of Senegal. Dorsal surface, scale bar = 100 mm.

Sl. 2: Normalni primerek vrste *Zanobatus schoenleinii* (Ref. ISPAB- Zan-sch-01) iz senegalske obale. Hrbtna stran, merilo = 100 mm.

morphological characters: disc sub-circular, wider than long; snout blunt, angle nearly 120° in front; nostrils narrow, anterior valves connected across the internarial space; mouth straight; teeth small; spiracles large without folds; dorsal and caudal fins small and rounded; covered by a rigid skin, unlike and minute scales, a medial row of thorns in disk and tail, and three rows arranged in an arc of circle on each shoulder; back brown with dark cross bands with white spots in between near the pectoral edges (Tab. 1; Fig. 2). Such description is in total agreement with Garman (1913), Cadenat (1951), Blache et al. (1970), and Séret & Opic (1990).

The abnormal specimen was a sub-adult male of *Zanobatus schoenleinii*, with developed but flexible claspers, measuring 495 mm TL, 279 mm DW and 751 g TBW (Tab. 1; Fig. 3). The central part of the disc was brown with dark spots externally larger and abundant on the central part of the disc, conversely, dark cross bands were totally lacking. An area surrounding eyes and spiracles was clear and whitish. The edges of both pectorals were largely clear, both dorsal fins and caudal fin whitish, belly entirely beige. The eye colour was black, which confirmed that the specimen was simultaneously leucistic and

piebald and not a true albino (*sensu* Quigley et al. (2019)). It is the second case reported for *Z. schoenleinii*, a similar specimen was previously collected from the coast of Senegal (Diatta et al., 2013).

Quigley et al. (2019) noted that albinism is a disorder controlled by different genes in which the pigment melanin is absent or non-functional. Additionally, leucism is a partial lack of pigmentation of the skin controlled by a single recessive allele (Jones et al., 2016). The term partial albinism or pseudo-albinism was previously used, but is now considered obsolete (Van Grouw, 2006). Lipej et al. (2011) showed that large albino specimens of marbled torpedo *Torpedo marmorata* Risso, 1810 could also be captured in the wild and that, consequently, total or partial albinism does not compromise life and development. Similar patterns were also reported for the striped panray (Diatta et al., 2013). The relationship log DW versus log TBW is $\log \text{TBW} = -6.97 + 4.07 * \log \text{DW}$; $r = 0.94$, $n = 52$. It appears a positive allometry but the leucistic piebald specimen which clearly displays a TBW lower than this of the other specimens (Fig. 4). Following the opinion of Quigley et al. (2019), such pattern suggests that this abnormal specimen did not develop like other normal specimens in the wild due to being




Fig. 3: Leucistic piebald specimen of *Zanobatus schoenleinii* (Ref. ISPAB- Zan-sch-01) from the coast of Senegal. A. Dorsal surface, scale bar = 80 mm. B. Ventral surface, scale bar = 100 mm.

Sl. 3: Belični piebalistični primerek vrste *Zanobatus schoenleinii* (Ref. ISPAB- Zan-sch-01) iz senegalske obale. A. Hrbtna stran, merilo = 80 mm. B. Trebušna stran, merilo = 100 mm.

Tab. 1: The morphometric measurements (in mm and as % DW) and total body weight recorded in the leucistic piebald specimen of *Zanobatus schoenleinii* (ISPAB-Zan-sch-02) and compared to a normal specimen (ISPAB-Zan-sch-01), both from the coast of Senegal.

Tab. 1: Morfometrične meritve (v mm in kot % DW) in celotna telesna masa pri beličnemu piebaldističnemu primerku vrste *Zanobatus schoenleinii* (ISPAB-Zan-sch-02) v primerjavi z normalnim primerkom (ISPAB-Zan-sch-01). Oba sta bila ujeta ob senegalski obali.

References	ISPAB-Zan-sch-01		ISPAB-Zan-sch-02	
Measurements	MM	% DW	MM	% DW
Total length	559	194.8	495	177.4
Disc length	313	94.8	255	91.4
Disc width	330	100.0	279	100.0
Disc depth	44	13.3	34	12.2
Eyeball length	17	5.2	18	6.5
Pre-orbital length	70	21.2	59	21.2
Inter-orbital length	26	7.9	28	10.0
Spiracle diameter	23	6.7	19	6.8
Interspiracular width	33	10.0	30	10.8
Space between eye and spiracle	13	3.9	11	3.9
Pre-oral length	72	21.2	60	21.5
Mouth width	43	13.0	43	15.4
First gill-slit	8	2.5	9	3.2
Fifth gill-slit	7	2.4	7	2.5
Width between first gill-slit	82	24.8	69	24.7
Width between fifth gill-slit	51	15.5	43	15.4
Snout tip to eye	81	24.3	68	24.4
Snout tip to mouth	81	24.3	68	24.4
Snout tip to first gill-slit	106	32.1	102	36.6
Snout tip to fifth gill-slit	150	45.5	132	47.3
Snout tip to vent	270	81.8	225	80.7
Pectoral fin anterior margin	205	71.4	185	66.3
Pectoral fin posterior margin	184	62.1	143	51.3
Pelvic fin anterior margin	50	15.2	48	17.2
Pelvic fin posterior margin	77	23.3	55	19.7
Second dorsal inneredge	10	3.0	11	3.9
Interdorsal distance	30	9.1	32	11.5
Second dorsal to caudal birth	28	8.6	35	12.5
Total body weight	1507		751	

maladapted. But it could also explain why of the 633 batoid species known to date worldwide, only 22 cases of albinism, leucism and piebaldisim had been reported (Quigley et al., 2018), 23 including the present specimen.

Talent (1973) noted that lack of pigmentation could be considered as disadvantage for sharks and rays, calling the attention of possible predators to their presence in the wild. Generally, abnormal specimens are considered of lower value than normal specimens among fishermen and are sometimes discarded at sea; this could also explain their scarcity in fish landings. Unfavourable environmental conditions, such as large exposure to pollutants, could also play a role in the lack of pigmentation, especially in species living a benthic zones, such as *Zanobatus schoenleinii*. Such phenomenon was particularly observed in batoids, which generally inhabit sandy-muddy bottoms (Ribeiro-Prado et al., 2008; Diatta et al., 2013). The fact that albinos, leucistic and piebald specimens are more sensitive to pollutants than normal specimens remains a valid hypothesis, which should, however, be validated by further studies. Since Diop et al. (2012) and Bonnin et al. (2016) report that the coast of Senegal, especially around the touristic area of Dakar, has been, for decades, increasingly




Fig. 4: Total body mass (TBM) versus disc width (DW) relationship expressed in logarithmic co-ordinates for normal and leucistic piebald specimens of *Zanobatus schoenleinii* collected from the coast of Senegal.
Sl. 4: Odnos med celotno telesno maso (TBM) in širino diska (DW), izražen v logaritmični skali pri normalnih in leucističnih piebaldističnih primerkih vrste *Zanobatus schoenleinii*, ujetih ob senegalski obali.

exposed to pollutants in the wild and their impact on the local biodiversity cannot be totally ruled out, the presently studied case could be a good instance of that.

BELIČNI PRIMEREK VRSTE *ZANOBATUS SCHOENLEINII* (CHONDRICHTHYES: ZANOBATIDAE) IZ SENEGALSKE OBALE (VZHODNI TROPSKI ATLANTIK)

Christian CAPAPÉ

Laboratoire d'Ichtyologie, Université de Montpellier, case 104, 34095 Montpellier cedex 5, France
e-mail: capape@univ-montp2.fr

Youssouph DIATTA & Almamy DIABY

Laboratoire de Biologie marine, Institut fondamental d'Afrique noire, (IFAN Ch. A. Diop), Université Cheikh Anta Diop de Dakar,
BP 206, Dakar, Senegal

Sihem RAFRAFI-NOUIRA

Université de Carthage, Unité de Recherches Exploitation des Milieux aquatiques, Institut Supérieur de Pêche et d'Aquaculture de Bizerte,
BP 15, 7080 Menzel Jemil, Tunisia

Christian REYNAUD

Laboratoire Interdisciplinaire en Didactique, Education et Formation, Université de Montpellier, 2 place Marcel Godechot, B.P. 4152,
34092 Montpellier cedex 5, France

ABSTRACT

V pričujočem prispevku avtorji poročajo o primeru beličnosti/piebaldzima pri vrsti skata *Zanobatus schoenleinii* (Müller & Henle, 1841), ujetega v plitvini ob obali Senegala (tropski vzhodni Atlantik). Gre za drugi primer pojava te anomalije pri tej vrsti. Primerek je bil subadultni samec, ki je meril 495 mm v dolžino, 279 mm v premeru diska in tehtal 751 g, ter imel razvita, a še vedno prožna klasperja. Avtorji nadalje razpravljajo o primerih albinizma, še posebej tistih pri skatih.

Ključne besede: morfologija, barva, pigmentacija, morfometrične meritve, skati

REFERENCES

- Blache J., J. Cadenat & J. Stauch (1970):** Clés de détermination des poissons de mer signalés dans l'Atlantique oriental tropical (entre le 20 ème parallèle N. et le 15 ème parallèle S. Faune trop. ORSTOM, 18, 1-479.
- Bonnin, M., I. Ly, B. Queffelec & M. Ngaido (2016):** Droit de l'environnement marin et côtier au Sénégal, IRD, PRCM, Dakar, Sénégal, 532 pp.
- Cadenat, J. (1951):** Poissons de mer du Sénégal. Init. Afr. Inst. Fr. Afr. Noire, Dakar, 3, 1-345.
- Capapé, C., M. N'Dao & M. Diop (1995):** Données sur la biologie de la reproduction de quatorze espèces de Sélaçiens batoïdes capturés dans la région marine de Dakar-Ouakam (Sénégal, Atlantique oriental tropical). Bull. Inst. fond. Afr. noire Cheikh Anta Diop, Dakar, sér. A, 48, 89-102.
- Diatta, Y., C. Reynaud & C. Capapé (2013):** First case of albinism recorded in striped panray, *Zanobatus schoenleinii* (Chondrichthyes: Platyrhinidae) from the coast of Senegal (Eastern Tropical Atlantic). J. Ichthyol., 53(11), 1007-1012.
- Diop, C., D. Dewaele, A. Toure, M. Cabral, F. Cazier, M. Fall, B. Ouddane & A. Diouf (2012):** Study of sediment contamination by trace metals at wastewater discharge points in Dakar (Senegal). J. Wat. Sci., 25(3), 185-299.
- Froese, R., A.C. Tsikliras & K.I. Stergiou (2011):** Editorial note on weight-length relations of fishes. Acta Ichthyol. Piscat., 41(4), 261-263.
- Garman, S. (1913):** The Plagiostoma (Sharks, Skates, and Rays). Mem. Mus. Comp. Zool. Harvard Univ., 36(2), 1-515.
- Jones, C.M., E.R. Hoffmayer & R.P. Gropp (2016):** First record of a leucistic *Narcine bancrofti* (Elasmobranchii, Narcinidae) from the northern Gulf of Mexico. Cybium, 40(3), 249-251.
- Lipej, L., B. Mavrič, V. Žiža & C. Capapé (2011):** First cases of albinism recorded in the marble electric ray *Torpedo marmorata* (Chondrichthyes: Torpedinidae). Cah. Biol. Mar., 52(3), 261-267.
- Lloris, D. & J. Rocabado (1998):** Guide FAO d'identification des espèces pour les besoins de la pêche. Guide d'identification des ressources marines vivantes pour le Maroc. FAO, Rome, 263pp.
- Maurin, C. & M. Bonnet (1970):** Poissons des côtes nord-ouest africaines (Campagnes de la «Thalassa», 1962 et 1968). Rev. Trav. Inst. scient. Tech. Pêch. marit., 34, 125-170.
- Pauly, D. (1983):** Some simple methods for assessment of tropical fishes. FAO Fish. Techn. Pap., 234: 3-10.

Quigley, D.T.G., A. de Carlos, D. Barros-Garcia & D. McGabhan (2018): Albinism and leucism in blonde ray (*Raja brachyura* Lafont, 1871) (Elasmobranchii: Batoidea) from the Irish Sea. Bull. Eur. Ass. Fish Pathol., 38(2), 79-88.

Quigley, D.T.G., A. de Carlos, D. Barros-Garcia & D. McGabhan (2019): Leucistic piebald Cuckoo ray (*Leucoraja naevus*) (Müller and Henle) from the Irish Sea. Irish Natur.J., 36(2), 159-162.

Ribeiro-Prado, C.C., M.C. Oddone, M.M. Bueno Gonzalez, A. Ferreira de Amorim & C. Capapé (2008): Morphological Abnormalities in Skates and Rays (Chondrichthyes) from off Southeastern Brazil. Arq. Cienc. Mar. Fortaleza, 41(2), 21-28.

Sanches, J.G. (1991): Catálogo dos principais peixes marinhos da República da Guiné-Bissau. Publicações avulsas do Instituto Nacional de Investigação das Pescas, Lisboa, 16, 1-429.

Séret, B. & P. Opic (1990): Poissons de mer de l'ouest Africain tropical. Initiations-Documentations Techniques, ORSTOM: Paris, 450 pp.

Talent, L.G. (1973): Albinism in embryo gray smoothhound sharks, *Mustelus californicus*, from Elkhorn Slough, Monterey Bay, California. Copeia, 3, 595-597.

Van Grouw, H. (2006): Not every white is an albino: sense and nonsense about colour aberration in birds. Dutch Birding, 28(3), 79-89.