

ANNALES



*Analí za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 30, 2020, 2*



UDK 5

ISSN 1408-533X
e-ISSN 2591-1783



ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 30, 2020, 2

KOPER 2020

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitzsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Lektor/Supervisione/Language editor:

Petricja Berlot Kužner (angl.)

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelja/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria[®]Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente[®]**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 21. 12. 2020.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS), Luka Koper in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation: 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: Clarivate Analytics: Biological Abstracts / BIOSIS Previews / Zoological Record; Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
 The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
 Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / INDICE GENERALE / CONTENTS 2020(2)

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
MEDITERRANEAN SHARKS**Paraskevi K. KARACHLE, Caterina STAMOULI & Aikaterini DOGRAMMATI**

Review of the Sharpnose Sevengill Shark
Heptranchias perlo (Chondrichthyes:
Hexanchidae) in the Mediterranean:
Historical and Recent Data 131
Pregled o pojavljanju morskega psa
sedmoroškrgarja Heptranchias perlo
(Chondrichthyes: Hexanchidae) v Sredozemlju:
zgodovinski in recentni podatki

Hakan KABASAKAL

A Historical Catch of White Shark,
Carcharodon carcharias (Lamniformes:
Lamnidae), in the Sea of Marmara (Turkey)
from the 1950s 147
Zgodovinski pregled ulova belega morskega
volka, Carcharodon carcharias (Lamniformes:
Lamnidae), v Marmarskem morju (Turčija) iz
petdesetih let

Nexhip HYSOLAKOJ, Rigers BAKIU & Alen SOLDO

The First Record of Kitefin Shark
Dalatias licha in Albanian Waters 151
Prvi zapis o pojavljanju klinoplavutega
morskega psa Dalatias licha v albanskih vodah

Deniz ERGÜDEN, Mahmut İĞDE, Cemal TURAN, Deniz AYAS & Hakan KABASAKAL

Occurrence of a Large Bigeye Thresher Shark,
Alopias superciliosus (Lamniformes: Alopiidae),
in the Northeastern Levantine Sea (İskenderun
Bay, Eastern Mediterranean Sea, Turkey) 157
Pojavljanje velike velikooke morske lisice,
Alopias superciliosus (Lamniformes: Alopiidae),
v severovzhodnem Levantskem morju (zaliv
İskenderun, vzhodni Mediteran, Turčija)

Lovrenc LIPEJ, Janja FRANCÉ, Domen TRKOV, Borut MAVRIČ & Aleš BOLJE

The Occurrence and Status of Thresher Shark
(Alopias vulpinus) in Waters Off Slovenia 165
Pojavljanje in status navadnih morskih lisic
(*Alopias vulpinus*) v vodah Slovenije

Hakan KABASAKAL & Erdi BAYRI

First Record of a Young-Of-The-Year
Carcharodon carcharias in the
Strait of the Dardanelles 175
Prvi zapis o pojavljanju prvoletnega
belega morskega volka (*Carcharodon*
carcharias) v dardanski ozini

Francesco TIRALONGO, Clara MONACO & Alessandro DE MADDALENA

Report on a Great White Shark *Carcharodon*
carcharias Observed Off Lampedusa, Italy 181
Zapis o pojavljanju belega morskega volka
Carcharodon carcharias pri Lampedusi, Italija

Hakan KABASAKAL

A Leucistic White Shark, *Carcharodon*
carcharias (Lamniformes: Lamnidae),
from the Northern Aegean Sea, Turkey 187
Belični primerek belega morskega volka,
Carcharodon carcharias (Lamniformes:
Lamnidae), iz severnega Egejskega
morja, Turčija

IHTIOLOGIJA

ITTOLOGIA
ICHTHYOLOGY**Christian CAPAPÉ, Youssouph DIATTA, Almamy DIABY, Sihem RAFRAFI-NOUIRA & Christian REYNAUD**

Leucistic Piebald Striped Panray,
Zanobatus schoenleinii (Chondrichthyes:
Zanobatidae), from the Coast of Senegal
(Eastern Tropical Atlantic) 193
Belični primerek vrste Zanobatus
schoenleinii (Chondrichthyes:
Zanobatidae) iz Senegalske obale
(vzhodni tropski Atlantik)

Fatiha DALOUCHÉ, Lotfi BENSAHLA-TALET, Sidi Mohamed El Amine ABI AYAD & Ahmed BENSAHLA-TALET

Gonadal Development of Bogues,
Boops boops (Linnaeus, 1758),
from Oran Bay (Teleostei, Sparidae) 201
Razvoj gonad pri bukvi, *Boops boops*
(Linnaeus, 1758), iz Oranskega zaliva
(Teleostei, Sparidae)

Francesco TIRALONGO & Roberto PILLON	FLORA
New Distributional Records of <i>Gobius buccichi</i> (Pisces, Gobiidae) from the Mediterranean Sea and In Situ Comparisons with <i>Gobius incognitus</i> 215	FLORA
<i>Novi podatki o razširjenosti vrste glavača Gobius buccichi (Pisces, Gobiidae) iz Sredozemskega morja in in situ primerjave z vrsto Gobius incognitus</i>	FLORA
BIOINVAZIJA	Amelio PEZZETTA
BIOINVASIONE	Le Papaveraceae Juss. della Flora Italiana: Distribuzione Regionale e Considerazioni Fitogeografiche 245
BIOINVASION	<i>Družina Papaveraceae Juss. v italijanski flori: regionalna razširjenost in fitogeografska opredelitev</i>
İnci TÜNEY-KIZILKAYA & Okan AKYOL	Toufik CHEDADI, Omar IDRISI, Anas ELKHABLI, Youssef KHACHTIB, Abdelmajid HADDIOUI & Mohammed EL HANSALI
Occurrence of <i>Scarus ghobban</i> (Scaridae) at the Border of the Aegean Sea (Kaş, Turkey) 223	Morphological Characterization of <i>Orobanche crenata</i> in Carrots and Legumes (Faba Bean and Chickpea): Indications of Potential Genetic Differentiation Towards Host Plants 265
<i>Pojavljanje modroprogaste morske papige (Scaridae) na meji Egejskega morja (Kaş, Turkey)</i>	<i>Morfološka opredelitev vrste Orobanche crenata pri korenju in stročnicah (bob in čičerika): pokazatelji možne genetske diferenciacije proti gostiteljskim rastlinam</i>
Argyro ZENETOS & Anastasia MILIOU	OCENE IN POROČILA
Abudefduf cf. <i>saxatilis</i> in the Saronikos Gulf, Greece: Unaided Introduction or Human Aided Transfer? 227	RECENSIONI E RELAZIONI
Abudefduf cf. <i>saxatilis</i> v zalivu Saronikos (Grčija): nenameren ali nameren vnos?	REVIEWS AND REPORTS
JADRANSKA MORSKA BIODIVERZITETA	Lovrenc Lipej
BIODIVERSITÀ MARINA DELL'ADRIATICO	Book review: ICHTHYOFAUNA OF THE ADRIATIC SEA 283
ADRIATIC MARINE BIODIVERSITY	Lovrenc Lipej
Martina ORLANDO-BONACA & Domen TRKOV	Book review: KNIGHTS, BALLERINAS AND INVISIBLES: THE DECAPOD CRUSTACEANS OF THE BRIJUNI MARINE PROTECTED AREA 284
After More Than Forty-Five Years a New Finding of <i>Cystoseira foeniculacea</i> f. <i>latiramosa</i> in the Coastal Sea of Slovenia 233	Alenka Malej
<i>Nova najdba vrste Cystoseira foeniculacea f. latiramosa v obalnem morju Slovenije po več kot štiridesetih letih</i>	Book review: BIODIVERZITETA BIOGENIH FORMACIJ. ZAKLADNICA NARAVE SLOVENSKEGA MORJA 285
Valentina PITACCO	POPRAVKI IN DOPOLNILA
First Record of the Echiurid <i>Maxmuelleria gigas</i> (M. Müller, 1852) in Slovenian Waters (Northern Adriatic) 239	ERRATA CORRIGE
<i>Prvi zapis o pojavljanju zvezdaša Maxmuelleria gigas (M. Müller, 1852) v slovenskih vodah (severni Jadran)</i>	ERRATA & CORRIGENDA
	Corrigendum to “Capture of a Large Great White Shark, <i>Carcharodon carcharias</i> (Lamnidae) from the Tunisian Coast (Central Mediterranean Sea): a Historical and Ichthyological Event”, Annales, Series Historia Naturalis, 2020, 30(1), 9-14 289
	Kazalo k slikam na ovitku 290
	<i>Index to images on the cover</i> 290

received: 2020-06-10

DOI 10.19233/ASHN.2020.18

OCCURRENCE OF A LARGE BIGEYE THRESHER SHARK, *ALOPIAS SUPERCILIOSUS* (LAMNIFORMES: ALOPIIDAE), IN THE NORTHEASTERN LEVANTINE SEA (ISKENDERUN BAY, EASTERN MEDITERRANEAN SEA, TURKEY)

Deniz ERGÜDEN, Mahmut İĞDE & Cemal TURAN

İskenderun Technical University, Marine Sciences and Technology Faculty, Marine Sciences Department, İskenderun, Hatay, Turkey
e-mail: deniz.erguden@iste.edu.tr

Deniz AYAS

Faculty of Fisheries, Mersin University Yenişehir Campus, 33160, Mersin, Turkey

Hakan KABASAKAL

Ichthyological Research Society, Tantavi mahallesi, Menteşoğlu caddesi, İdil apartmanı, No: 30, D: 4, 34764 Ümraniye, İstanbul, Turkey
e-mail: kabasakal.hakan@gmail.com

ABSTRACT

On 9 April 2019, a single female specimen of *Alopias superciliosus* Lowe, 1841 was incidentally captured by a commercial purse seiner at an approximate depth of 20 m, off the coast of Çevlik, İskenderun Bay, Turkey. The present specimen is the largest to be captured in northeastern Levantine Sea to date, and one of the largest specimens, approaching the alleged 500 cm maximum TL, caught in the entire Mediterranean Sea. While the congeneric thresher shark, *A. vulpinus*, is now listed as protected species in the Turkish Marine Fisheries Act, a similar conservation status for the protection of *A. superciliosus* is urgently needed.

Key words: thresher shark, Alopiidae, coast of Cevlik, Levantine Sea, conservation, bycatch

PRESenza DI SQUALO VOLPE OCCHIONE, *ALOPIAS SUPERCILIOSUS* (LAMNIFORMES: ALOPIIDAE), NEL MAR LEVANTINO NORD-ORIENTALE (BAIA DI ISKENDERUN, MEDITERRANEO ORIENTALE, TURCHIA)

SINTESI

Il 9 aprile 2019, una femmina di squalo volpe occhione, *Alopias superciliosus* Lowe, 1841, è stata accidentalmente catturata da una rete a ciruizione commerciale, a una profondità di circa 20 m, al largo della costa di Çevlik, nella baia di İskenderun, in Turchia. L'esemplare è il più grande ad essere stato catturato nel Mar Levantino nord-orientale fino ad oggi. Si tratta inoltre di uno degli esemplari più grandi, avvicinandosi al presunto limite massimo di 500 cm, catturato in tutto il mare Mediterraneo. Mentre lo squalo volpe, *A. vulpinus*, è ora considerato specie protetta dalla legge turca sulla pesca marittima, è urgentemente necessario uno stato di conservazione simile per la protezione di *A. superciliosus*.

Parole chiave: squalo volpe occhione, Alopiidae, costa di Cevlik, Mar Levantino, conservazione, catture accidentali

INTRODUCTION

The bigeye thresher shark, *Alopias superciliosus* Lowe, 1841, is one of the three species in the family Alopiidae, which is distributed worldwide in all temperate and tropical oceans (Ebert et al., 2013). Despite its occurrence in the epipelagic and mesopelagic zones of open oceans, *A. superciliosus* is a common inhabitant of coastal waters over continental shelves, with seldom occurrence in shallow inshore waters (Froese & Pauly, 2020).

The first record of the bigeye thresher shark in the Mediterranean was reported from Italian waters, during observations in the fishing centres of Mazara del Vallo (Trapani, Sicily) (Cigala-Fulgosi, 1983), and was followed by a few dozen further records from elsewhere within the species' distribution range, which extends from the west to the east of the Mediterranean, including the Marmara Sea, which were summarised by Lanteri et al. (2017). *A. superciliosus* was recorded for the first time in Turkish waters by Mater (2005) in the Bay of Gökova.

This article reports on the capture of a large female of *A. superciliosus* in the Bay of İskenderun, north-eastern Levantine waters, and presents morphometric measurements of the examined specimen.

MATERIAL AND METHODS

On 9 April 2019, a single female specimen of *A. superciliosus* was incidentally captured by a commercial purse seiner at an approximate depth of 20 m, off the coast of Çevlik, İskenderun Bay, Turkey (location: 35° 57' 14" N, 35° 54' 30" E; Fig. 1). The specimen was photographed (Figs. 2, 3 and 4), and morphometric measurements (in cm) and weight (in grams) were recorded on site. Morphometric measurements were based on Compagno (2001), and measured to the nearest millimetre. Identification of the examined species follows Compagno (2001) and Serena (2005), and taxonomic nomenclature follows Froese & Pauly (2020).

RESULTS AND DISCUSSION

The total length (TL) of the examined bigeye thresher shark (Figs. 2, 3 and 4) was 472 cm, and its weight 400 kg. The morphometric measurements of the examined female are provided in Table 1, together with morphometric data extracted from several articles on *A. superciliosus*. Based on the following descriptive features that coincided with those proposed in Compagno (2001) and Serena (2005), the



Fig. 1: Map depicting the approximate site of capture (*) of the examined *A. superciliosus*.

Sl. 1: Zemljevid obravnavanega območja s približno označeno lokaliteto (*), kjer je bil analiziran primerek vrste *A. superciliosus* ujet.

Tab. 1: Comparison of selected morphometric measurements of the *A. superciliatus* specimen examined in this study with those of other reported specimens from the Mediterranean Sea. Values provided in parentheses are the percentages that selected morphometric measurement take up of TL. *Since Lanteri et al. (2017) base their record on the collected head and pectoral fins of the examined specimen, the TL value is not provided. W Med: western Mediterranean; NW Med: northwestern Mediterranean; SE Med: southeastern Mediterranean; NE Med: northeastern Mediterranean; N/A: not available.

Tab. 1: Primerjava izbranih morfometričnih meritev na pregledanem primerku vrste *A. superciliatus* v tej raziskavi s podatki, pridobljenimi iz drugih raziskav v Sredozemskem morju. Vrednosti v oklepaju so deleži izbranih morfometričnih meritev glede na celotno dolžino telesa. *Ker so Lanteri in sod. (2017) zbrali le glavo in prsne plavuti pregledanega primerka, dolžina telesa ni navedena. W Med: zahodno Sredozemlje; NW Med: severozahodno Sredozemlje; SE Med: jugovzhodno Sredozemlje; NE Med: severovzhodno Sredozemlje; N/A: ni podatka.

	This study	Kabasakal et. al. (2011)	Kabasakal (2017)	Ayas et al. (2020)	Corsini-Foka & Sioulas (2009)	Lanteri et al. (2017)*	Farrag (2017)
Locality	İskenderun Bay (SE Med. Turkey)	Fethiye Coast (Aegean Sea, Turkey)	Antalya Bay (W Med. Turkey)	Mersin Bay (NE Med. Turkey)	Dodecanese waters (Aegean Sea, Greece)	Ligurian Sea (NW Med. Italy)	Egyptian coast of Med. Sea, Egypt
Sex	Female	Female	Female	Female	Male	Female	N/A
Measurements	Value (cm) (% of TL)	Value (cm) (% of TL)	Value (cm) (% of TL)	Value (cm) (% of TL)	Value (cm) (% of TL)	Value (cm)	Value (cm) (% of TL)
Total Length (TL)	472.0	450.0	342.4	240.0	310.0	N/A	180.0
First dorsal fin length	40.0	41.0 (9.1%)	-	17.9 (7.4%)	27.0 (8.7%)	-	-
First dorsal fin base length	32.0 (6.8%)	32.0 (7.1%)	-	13.6 (5.6%)	21.0 (6.8%)	-	-
Pectoral fin length	83.0 (17.5%)	82.0 (18.2%)	-	51.5 (21.4%)	62 (20.0%)	25.4	35 (19.4%)
Pectoral fin base length	40.0 (8.5%)	30.0 (6.7%)	-	20.1 (8.4%)	21 (6.8%)	-	-
Ventral fin length	37.0 (7.8%)	39.0 (8.7%)	-	20.6 (8.6%)	26.0 (8.4%)	-	-
Ventral fin base length	30.0 (6.4%)	30.0 (6.7%)	-	15.4 (6.4%)	21.0 (6.8%)	-	-
Distance between dorsal fin origin and ventral fin origin	-	-	-	17.8 (7.4%)	23.0 (7.4%)	-	-
Distance between pectoral fin origin and ventral fin origin	204 (43.2%)	-	-	68.9 (28.7%)	77.0 (24.8%)	-	-
Predorsal length	-	128.0 (28.4%)	-	87.7 (36.5%)	94.0 (30.3%)	-	57.0 (31.6%)
Preventral length	-	165.0 (36.7%)	-	105.3 (43.9%)	118.0 (38.1%)	-	70.0 (38.9%)
Prepectoral length	-	55.0 (12.2%)	38.6 (11.2%)	43.7 (18.2%)	45.0 (14.5%)	-	29.0 (16.1%)
Preanal length	-	-	-	-	-	41.7	89.0 (49.4%)
Precaudal fin length	-	-	240.5 (70.2%)	-	-	-	-
Prebranchial length	-	-	34.6 (10.1%)	-	-	-	-
Preorbital length	-	-	14.7 (4.2%)	-	-	31.6	9.0 (5.0%)
Preoral length	-	-	15.1 (4.4%)	-	-	11.0	-
Interorbital space	-	-	9.4 (2.7%)	-	-	12.3	7.0 (3.9%)
Eye length	-	-	6.9 (2.01%)	-	-	7.4	4.5 (0.95%)
Eye height	-	-	5.1 (1.4%)	-	-	4.7	-
Mouth width	20.0 (4.2%)	-	-	-	-	7.4	-
Pectoral fin anterior margin	-	-	46.4 (13.5%)	-	-	10.4	-
Tail length	240.0 (50.5%)	217.0 (48.2%)	-	103.8 (43.3%)	143.0 (46.1%)	54.4	84.0 (17.8%)




Fig. 2: Female *A. superciliatus* (472 cm in TL), captured off the coast of Çevlik, northeastern Levantine Sea (Photo: D. Ergüden archive).

Sl. 2: Samica vrste *A. superciliatus* (472 cm v dolžino telesa), ujeta ob obali Çevlika, severovzhodno Levantsko morje (Foto: D. Ergüden arhiv).

examined specimen was identified as *A. superciliatus*: cylindrical body, massive before the first dorsal fin; snout moderately long and bulbous; head with deep grooves extending along either sides from behind the eyes to the areas above the gill openings; eyes large and reaching the dorsal surface of head; interorbital space nearly flat; first dorsal midbase closer to the pelvic fin base than to the pectoral fin bases; pectoral fins falcate with broad apices; upper surface of the body brownish to dark grey, growing lighter in colour ventrally.

The story of *A. superciliatus* in the Mediterranean Sea dates to the early 1980s, when Cigala-Fulgosi (1983) recorded the species in the region for the first time. However, the discovery of taxidermied specimens caught in Dodecanese waters (southeastern Aegean Sea) in the 1950s suggests that *A. superciliatus* had been present in the Mediterranean Sea some thirty years earlier than the first published report on its occurrence was made (Corsini-Foka & Sioulas, 2009). Following Cigala-Fulgosi's (1983) report, several captures of bigeye thresher shark were reported from different parts of the Mediterranean Sea. These include Pescara, Italy (Cugini & De Maddalena, 2003), waters around Toulon (Capape, 1977), the Catalan littoral (Barrull et al., 1999), eastern Sardinia (Vacchi & Serena, 2000), and the Ligurian Sea (Lanteri et al., 2017). In the eastern Mediterranean region, sporadic records of *A. superciliatus* were reported from the Turkish Mediterranean and Aegean coasts (Mater, 2005; Kabasakal et al., 2011; Gökoğlu et al., 2017; Kabasakal, 2017; Ayas et al., 2020), as well as from the Sea of Marmara (Kabasakal & Karhan, 2008), Egyptian waters (Farrag, 2017), Israel, off the Ashod coast (Golani, 1996), Dodecanese waters, the SE Aegean Sea (Corsini-Foka & Sioulas, 2008), and from Cyprus waters (Kleitou et al., 2017). Lanteri et al. (2017) compiled a full list of 40 Mediterranean records of *A. superciliatus* with some data on the captured specimens and relevant references. With the addition of the specimens reported by Kabasakal (2017), Ayas et al. (2020), and the one recorded in the present study, the total number of records of *A. superciliatus* from the Mediterranean and Turkey increased to 43 and 15, respectively, with the latter number representing almost 35 % of all records reported in the Mediterranean to date. Although recent data suggest that the distribution of *A. superciliatus* in the Mediterranean is extending from the western part of the region to the Levantine Sea in the east and the Marmara Sea in the northernmost part, this shark species is still considered rare in the Mediterranean (Serena, 2005; Lanteri et al., 2017), and the small sample of 43 recorded specimens from the 1950s to date seems to validate this view. An overview of *A. superciliatus* records from Turkish waters is presented in Table 2.

While the maximum total length reported for *A. superciliatus* is 484 cm (Ebert & Stehmann, 2013), Lanteri et al. (2017) provide records on specimens that reach 500 cm. The specimen presented in our article is the largest ever caught in the northeastern Levantine Sea to date. It is also one of the largest specimens, approaching the alleged 500 cm maximum TL, that have ever been caught in the entire Mediterranean Sea. According to Ebert and Stehmann (2013), females mature between 332 and 356 cm TL; thus, the present specimen is considered a mature female.

Tab. 2: An overview of the records of *A. superciliosus* from Turkish waters. GN: gill net; SN: shrimp net; PS: purse seine; TN: trammel net; SN: stationary net; BT: bottom trawl; LL: longline; PSLL: pelagic swordfish longline.

Tab. 2: Pregled zapisov o pojavljanju vrste *A. superciliosus* v turških vodah. GN: zabodna mreža; SN: mreža za kozice; PS: zaporna plavarica; TN: trislojna mreža; SN: stoječa mreža; BT: pridnena povlečna mreža; LL: parangal; PSLL: parangal za mečarice.

No	Authors	Date	Depth (m)	Locality	Region	Gear	Sex	Length (cm, TL)	Weight (Kg)
1	Clo et al. (2008)	April 2005	-	Marmaris	SE Aegean Sea	GN	-	350	160
2	Mater (2005)	23 May 2005	12	Gökova	SE Aegean Sea	SN	-	350	150
3	Kabasakal & Karhan (2008)	25 February 2007	-	Silivri	Marmara Sea	PS	-	450	-
4	Kabasakal et al. (2011)	28 February 2011	110	Fethiye	SE Aegean Sea	TN	♀	430	300
5	Kabasakal et al. (2011)	2 July 2011	-	Silivri	Marmara Sea	PS	♀	250	65
6	Lanteri et al. (2017)	25 February 2013	-	-	Marmara Sea	-	-	370	250
7	Lanteri et al. (2017)	13 March 2013	-	Yeşilköy	Marmara Sea	-	-	200	110
8	Lanteri et al. (2017)	15 July 2013	-	Yediburunlar	SE Aegean Sea	-	-	500	95
9	Lanteri et al. (2017)	15 April 2015	-	Antalya	Mediterranean Sea	-	-	-	-
10	Kabasakal et al. (2011)	21 May 2016	100	Sivrice	NE Aegean Sea	SN	♀	400	-
11-12	Gökoglu et al. (2017)	March, April and July 2015	600-700	Antalya	Mediterranean Sea	BT, LL	♀, ♂	180-299	15.5-65
13	Kabasakal (2017)	22 December 2016	-	Antalya Bay	Mediterranean Sea	PSLL	♀	342.4	-
14	Ayas et al. (2020)	02 January 2020	25	Taşucu	Mediterranean Sea	TN	♀	240	48
15	Present specimen	09 April 2019	20	Keldağ, Çevlik Coast	Mediterranean Sea	PS	♀	472	400



Fig. 3: Front (A) and lateral (B) views of the examined *A. superciliosus* (Photo: D. Ergüden archive).

Sl. 3: Pogled od spredaj (A) in od strani (B) pregledanega primerka vrste *A. superciliosus* (Foto: D. Ergüden arhiv).



Fig. 4: Dentition on the upper and lower jaws of the examined *A. superciliatus* (Photo: D. Ergüden archive).
Sl. 4: Zobovje na spodnji in spodnji čeljustnici pri pregledanem primerku vrste *A. superciliatus* (Foto: D. Ergüden arhiv).

In the Mediterranean Sea, the bigeye thresher shark is considered occasional bycatch affected by interaction with pelagic fishing gear, as well as bottom trawls, longlines and entangling nets (Serena, 2005; Bariche, 2012). The previous IUCN category – “data deficient” - for *A. superciliatus* was recently changed to “endangered” for the Mediterranean Sea, and “vulnerable” worldwide (Otero et al., 2019). *A. superciliatus* is a typical *k*-selected species, with the maximum litter size ranging from 2 to 4 embryos (Bariche, 2012). While

the congeneric thresher shark, *A. vulpinus* is now listed as a protected species in the Turkish Marine Fisheries Act (Öztürk, 2018), a similar conservation status for the protection of *A. superciliatus* is urgently needed.

ACKNOWLEDGEMENTS

Authors thank the fishermen who kindly permitted the taking of photographs and recording of the measurements of the specimen described in the article.

POJAVLJANJE VELIKE VELIKOOKE MORSKE LISICE, *ALOPIAS SUPERCILIOSUS* (LAMNIFORMES: ALOPIIDAE), V SEVEROVZHODNEM LEVANTSKEM MORJU (ZALIV ISKENDERUN, VZHODNI MEDITERAN, TURČIJA)

Deniz ERCÜDEN, Mahmut İĞDE & Cemal TURAN

İskenderun Technical University, Marine Sciences and Technology Faculty, Marine Sciences Department, İskenderun, Hatay, Turkey
e-mail: deniz.erguden@iste.edu.tr

Deniz AYAS

Faculty of Fisheries, Mersin University Yenişehir Campus, 33160, Mersin, Turkey

Hakan KABASAKAL

Ichthyological Research Society, Tantavi mahallesi, Menteşoğlu caddesi, İdil apartmanı, No: 30, D: 4, 34764 Ümraniye, İstanbul, Turkey
e-mail: kabasakal.hakan@gmail.com

POVZETEK

Devetega aprila 2019 so ujeli samico velikooke morske lisice *Alopias superciliatus* Lowe, 1841 v zaporno plavarico, na globini približno 20 m ob Cevliku v zalivu İskenderun Bay, v Turčiji. Gre za doslej največji primerek te vrste v severovzhodnem Levantskem morju in nasploh v Sredozemskem morju, saj je meril okoli 500 cm v dolžino. Njena sorodnica iz istega rodu, navadna morska lisica, *A. vulpinus*, je danes v turškem morskem ribolovnem aktu zavarovana vrsta, zato je nujno čimprej urediti podoben ohranitveni status tudi za vrsto *A. superciliatus*.

Ključne besede: morska lisica, Alopiidae, obala pri Cevliku, Levantsko morje, ohranjanje, prilog

REFERENCES

- Ayas, D., H.D. Akbora & B.A. Çiçek (2020):** Occurrence of the young thresher shark *Alopias superciliatus* Lowe, 1841 (Lamniformes: Alopiidae) in the Northeastern Mediterranean Sea. Comm. J. Biol., 4, 28-31.
- Bariche, M. (2012):** Field Identification Guide to the Living Marine Resources of the Eastern and Southern Mediterranean. FAO Species Identification Guide for Fishery Purposes. Rome, FAO, 610 pp.
- Barrull, J., I. Mate & M. Bueno (1999):** Observaciones de tiburones (chondrichthyes Euselachii) en aguas de Cataluña (Mediterraneo No), con algunos aspectos generales de su ecología. Sci. Gerund., 24, 127-151.
- Capape, C. (1977):** Liste commentee des selaciens de la region de Toulon (de la ciotat a saint-Tropez). Bull. Mus. Hist. Nature, Marseille, 37, 5-9.
- Clo, S., R. Bonfil & E. de Sabata (2008):** Additional records of the bigeye thresher shark, *Alopias superciliatus*, from the central and eastern Mediterranean Sea. JMB2- Biodiversity Records 6168. <http://www.mba.ac.uk/jmba/jmba2biodiversityrecords.php>.
- Cigala-Fulgosi, F. (1983):** First record of *Alopias superciliatus* (Lowe, 1840) in the Mediterranean, with notes on some fossil species of the genus *Alopias*. Ann. Mus. Civ. Stor. Nat. Genova, 84, 211-229.
- Compagno, L.J.V. (2001):** Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Vol. 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). FAO Spec. Cat. Fish. Purp. 1(2): 269 p. FAO, Rome.
- Corsini-Foka, M. & A. Sioulas (2009):** On two old specimens of *Alopias superciliatus* (Chondrichthyes: Alopiidae) from the Aegean waters. Mar. Biodiversity Rec., 2: e72.
- Cugini, G. & A. De Maddalena (2003):** Sharks captured off Pescara (Italy, western Adriatic Sea). Annales Ser. His. Nat., 13, 201-208.
- Ebert, D.A., S. Fowler & L. J. V. Compagno (2013):** Sharks of the World. A Fully Illustrated Guide. Wild Nature Press, Plymouth, United Kingdom.
- Ebert, D.A. & M. Stehmann (2013):** Sharks, batoids and chimaeras of the North Atlantic. FAO Species Catalogue for Fishery Purposes. No. 7. FAO, Rome, 523 pp.
- Farrag, M.M.S. (2017):** New record of the bigeye thresher shark, *Alopias superciliatus* Lowe, 1841 (Family: Alopiidae) from the eastern Mediterranean Sea, Egypt. Int. J. Fish. Aquat. Stud., 5, 316-318.

- Froese, R. & D. Pauly (2020):** *Alopias superciliosus* Lowe, 1841 Bigeye thresher. Retrieved from <https://www.fishbase.se/summary/Alopiassuperciliosus.html>.
- Golani, D. (1996):** The marine ichthyofauna of the eastern Levant – history, inventory, and characterization. Israel J. Zool., 42, 15-55.
- Gökoğlu, M., S. Teker & D. Julian (2017):** First report of thresher sharks (Alopiidae) in the Gulf of Antalya. Iran. J. Fish. Sci., 16, 1108–1113.
- Kabasakal, H. & S.Ü. Karhan (2008):** On the occurrence of the bigeyethresher shark, *Alopias superciliosus* (Chondrichthyes: Alopiidae), in Turkish waters. Mar. Biodivers. Rec., 1, e69. <https://doi.org/10.1017/s1755267207007452>.
- Kabasakal, H., C. Dalyan & A. Yurtsever (2011):** Additional records of the Bigeye thresher shark *Alopias superciliosus* (Lowe, 1839) (Chondrichthyes: Lamniformes: Alopiidae) from Turkish waters. Annales Ser. Hist. Nat., 21, 143-148.
- Kabasakal, H. (2017):** Notes on historical and contemporary catches of lamniform sharks in Turkish waters. Annales Ser. Hist. Nat., 27, 51-58.
- Kabasakal, H., S.Ü. Karhan & S. Sakınan (2017):** Review of the distribution of large sharks in the seas of Turkey (Eastern Mediterranean). Cah. Biol. Mar., 58, 219-228. <https://doi.org/10.21411/CBM.A.96D9F948>.
- Kleitou, P.C., C.I. Antoniou, I. Giovos & D. Kletou (2017):** How accurately are we describing the longline bycatch? The case of the 'rare' shark *Alopias superciliosus* in eastern Mediterranean. Int. J. Fish. Aquat. Stud., 5, 375-378.
- Lanteri, L., L. Castellano & F. Garibaldi (2017):** New record of *Alopias superciliosus* Lowe, 1841 in the North-Western Mediterranean and annotated review of the Mediterranean records. Acta Adriat., 58, 313-324.
- Mater, S. (2005):** Denizlerimizde yeni bir köpekbalığı: *Alopias superciliosus* (Lowe, 1841). Aquaculture 1, 10: (in Turkish).
- Otero, M., F. Serena, V. Gerovasileiou, M. Barone, M. Bo, J.M. Arcos, A. Vulcano, & J. Xavier (2019):** Identification guide of vulnerable species incidentally caught in Mediterranean fisheries. IUCN, Malaga, Spain, 204 pp.
- Öztürk, B. (2018):** National action plan for the conservation of cartilaginous fishes in the Turkish water of the eastern Mediterranean Sea. J. Black Sea/Medit. Environ., 24, 91-96.
- Serena, F. (2005):** Field identification guide to the sharks and rays of the Mediterranean and Black Sea. FAO Species Identification Guide for Fishery Purposes. Rome: FAO. 97.
- Vacchi, M. & F. Serena (2000):** On a large specimen of bigeye thresher shark *Alopias superciliosus* (Lowe, 1839) (Chondrichthyes: Alopiidae) stranded in Tavolara Island (Eastern Sardinia, Mediterranean). Séret, B. and Sire, J.-Y. (eds). Proceedings of the 3rd European Elasmobranch Association Meeting, Boulogne-sur-Mer (France), 1999. Soc. Fr. Ichtyol and IRD Paris, 3, p. 84.