

# ANNALES



*Anali za istrske in mediteranske študije  
Annali di Studi istriani e mediterranei  
Annals for Istrian and Mediterranean Studies  
Series Historia Naturalis, 32, 2022, 1*



UDK 5

ISSN 1408-533X  
e-ISSN 2591-1783



# ANNALES

**Anali za istrske in mediteranske študije**  
**Annali di Studi istriani e mediterranei**  
**Annals for Istrian and Mediterranean Studies**

**Series Historia Naturalis, 32, 2022, 1**

KOPER 2022

**UREDNIŠKI ODBOR/  
COMITATO DI REDAZIONE/  
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnšič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitzsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/  
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/  
Redattore responsabile per le scienze  
naturali/Natural Science Editor:**

Lovrenc Lipej

**Urednica/Redattrice/Editor:**

Martina Orlando-Bonaca

**Prevajalci/Traduttori/Translators:**

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/  
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

**Tisk/Stampa/Print:**

Založništvo PADRE d.o.o.

**Izdajatelja/Editori/Published by:**

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria<sup>®</sup>

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente<sup>®</sup>

**Sedež uredništva/Sede della redazione/  
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;

**e-mail:** annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 30. 06. 2022.

**Sofinancirajo/Supporto finanziario/  
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/> / The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/> / Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



## VSEBINA / INDICE GENERALE / CONTENTS 2022(1)

SREDOZEMSKI MORSKI PSI  
SQUALI MEDITERRANEI  
MEDITERRANEAN SHARKS**Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ**

Observations on Thresher Shark, *Alopias vulpinus* (Chondrichthyes: Alopiidae) from the Coast of Algeria (Southwestern Mediterranean Sea) ..... 1  
*Opazovanja morskih lisic, Alopias vulpinus (Chondrichthyes: Alopiidae) ob alžirski obali (jugozahodno Sredozemsko morje)*

**Elif ÖZGÜR ÖZBEK & Hakan KABASAKAL**

Notes on Smoothback Angel Shark, *Squatina oculata* (Squatiniformes: Squatinidae) caught in the Gulf of Antalya ..... 9  
*Zapis o pegastih sklatih, Squatina oculata (Squatiniformes: Squatinidae), ujetih v Antalijskem zalivu*

**Alessandro PAGANO & Alessandro DE MADDALENA**

Underwater Observations of the Rare Angular Roughshark *Oxynotus centrina* (Chondrichthyes: Squalidae) in the Waters of Santa Tecla (Sicily, Italy) ..... 17  
*Podvodna opazovanja redkega morskega pršiča, Oxynotus centrina (Chondrichthyes: Squalidae) v vodah Sante Tecle (Sicilija, Italija)*

**Deniz ERGÜDEN, Deniz AYAS & Hakan KABASAKAL**

Morphometric Measurements of Three Young Carcharhinid Species from Northeastern Levant (Mediterranean Sea) ..... 25  
*Morfometrične meritve mladičev treh vrst morskih psov iz družine Carcharhinidae iz severnovzhodnega Levanta (Sredozemsko morje)*

**Hakan KABASAKAL**

Projections on the Future of Deep-Sea Sharks in the Sea of Marmara, Where Deep Zones Are Threatened by Deoxygenation: a Review ..... 35  
*Napovedi o prihodnosti globomorskih morskih psov v Marmarskem morju, ogroženem zaradi pomanjkanja kisika: pregled*

**BIOINVAZIJA**

**BIOINVASIONE**  
**BIOINVASION**

**Alan DEIDUN, Bruno ZAVA & Maria CORSINI-FOKA**

Distribution Extension of *Lutjanus argentimaculatus* (Lutjanidae) and *Psenes pellucidus* (Nomeidae) to the Waters of Malta, Central Mediterranean Sea ..... 49  
*Širjenje areala vrst Lutjanus argentimaculatus (Lutjanidae) in Psenes pellucidus (Nomeidae) v malteške vode (osrednje Sredozemsko morje)*

**Sami M. IBRAHIM, Abdulraziq A.**

**ABDULRRAZIQ, Abdulghani ABDULGHANI, Sara A.A. AL MABRUK, David SALVATORI, Bruno ZAVA, Maria CORSINI-FOKA & Alan DEIDUN**

First Record of *Enchelycore anatina* (Muraenidae) from Libyan Waters and an Additional Record from Southern Italy (Western Ionian Sea) ..... 59  
*Prvi zapis o pojavljanju kavljezobe murene Enchelycore anatina (Muraenidae) iz libijskih voda in dodatni zapis za južno Italijo (zahodno Jonsko morje)*

|   |     |   |     |
|---|-----|---|-----|
| <b>Rasha Ali HENEISH &amp; Samir Ibrahim RIZKALLA</b><br>Morphometric and Meristic Characteristics of a New Record of Bluespot Mullet <i>Crenimugil seheli</i> (Pisces: Mugilidae) in Egyptian Mediterranean waters .....   | 67  | <b>Deniz ERGÜDEN &amp; Cemal TURAN</b><br>A Rare Occurrence of <i>Carapus acus</i> (Carapidae) in the Eastern Mediterranean, Turkey .....<br><i>Redko pojavljanje strmorinca Carapus acus (Carapidae) v vzhodnem Sredozemskem morju (Turčija)</i>                                   | 113 |
| <i>Novi zapis o pojavljanju vrste Crenimugil seheli (Pisces: Mugilidae) v egiptovskih sredozemskih vodah in njene morfometrične in meristične značilnosti</i>   |     |   |     |
| <b>Yana SOLIMAN, Adib SAAD, Vienna HAMMOUD &amp; Christian CAPAPÉ</b><br>Heavy Metal Concentrations in Tissues of <i>Siganus rivulatus</i> (Siganidae) from the Syrian Coast (Eastern Mediterranean Sea) .....  | 75  | <b>Laith JAWAD, Murat ŞIRİN, Miloslav PETRTÝL, Ahmet ÖKTENER, Murat ÇELIK &amp; Audai QASIM</b><br>Skeletal Abnormalities in Four Fish Species Collected from the Sea of Marmara, Turkey .....  | 119 |
| <i>Vsebnost težkih kovin v tkivih marmoriranega morskega kunca Siganus rivulatus (Siganidae) iz sirske obale (vzhodno Sredozemsko morje)</i>  |     | <i>Skeletne anomalije pri štirih vrstah rib iz Marmarskega morja (Turčija)</i>  |     |
| <b>IHTIOLOGIJA<br/>ITTOLOGIA<br/>ICHTHYOLOGY</b>  |     |   |     |
| <b>Jihade ALAHYENE, Brahim CHIAHOU, Hammou EL HABOUZ &amp; Abdelbasset BEN-BANI</b><br>Length Based Growth Estimation of the Blue Shark <i>Prionace glauca</i> from the Moroccan Central Atlantic Coast .....   | 85  | <b>RAZMNÖŽEVALNA EKOLOGIJA<br/>ECOLOGIA RIPRODUTTIVA<br/>REPRODUCTIVE ECOLOGY</b>   |     |
| <i>Dolžinsko-masni odnos in ocena rasti pri sinjem morskem psu (Prionace glauca) iz osrednje atlantske obale Maroka</i>   |     |   |     |
| <b>Okan AKYOL, Altan LÖK &amp; Funda ERDEM</b><br>Occurrence of <i>Cubiceps gracilis</i> (Nomeidae) in the Eastern Mediterranean Sea .....  | 101 | <b>Amaria Latefa BOUZIANI, Khaled RAHMANI, Samira AIT DARNA, Alae Eddine BELMAHI, Sihem ABID KACHOUR &amp; Mohamed BOUDERBALA</b><br>Gonadal Histology in <i>Diplodus vulgaris</i> from the West Algerian Coast .....   | 137 |
| <i>Pojavljanje klateža, Cubiceps gracilis (Nomeidae), v vzhodnem Sredozemskem morju</i>   |     | <i>Histologija gonad pri navadnem šparu (Diplodus vulgaris) iz zahodne alžirske obale</i>   |     |
| <b>Farid HEMIDA, Boualem BRAHMI, Christian REYNAUD &amp; Christian CAPAPÉ</b><br>Occurrence of the Rare Driftfish <i>Cubiceps gracilis</i> (Nomeidae) from the Algerian Coast (Southwestern Mediterranean Sea) .....  | 107 | <b>Cheikhna Yero GANDEGA, Nassima EL OMRANI, Rezan O. RASHEED, Mohammed RAMDANI &amp; Roger FLOWER</b><br>The Growth and Reproduction of Two Sparidae, <i>Pagrus caeruleostictus</i> and <i>Pagellus bellottii</i> in Northern Mauritanian Waters (Eastern Tropical Atlantic) ..... | 143 |
| <i>Pojavljanje redkega klazeža Cubiceps gracilis (Nomeidae) z alžirske obale (jugozahodno Sredozemsko morje)</i>  |     | <i>Rast in razmnoževanje dveh vrst pagrov, Pagrus caeruleostictus in Pagellus bellottii v severnih mavretanskih vodah (vzhodni tropski Atlantik)</i>  |     |
| <b>Nassima EL OMRANI, Hammou EL HABO-UZ, Abdellah BOUHAIMI, Jaouad ABOU OUALID, Abdellatif MOUKRIM, Jamila GOZOULI, Mohammed RAMDANI, Roger FLOWER &amp; Abdelbasset BEN-BANI</b><br>The Reproductive Biology of the Pouting <i>Trisopterus luscus</i> from the Atlantic Coast of Morocco ..... |     |   |     |
|   |     | <i>Reproduktivna biologija francoskega moliča (Trisopterus luscus) iz atlantske obale Maroka</i>  | 155 |

**Mourad CHÉRIF, Rimel BENMESSAOUD & Christian CAPAPÉ**

- Growth Patterns and Age Structure of *Mullus surmuletus* (Mullidae) from the Northern Coast of Tunisia (Central Mediterranean Sea) ..... 173  
*Rastni parametri in starostna struktura progasti bradačev Mullus surmuletus (Mullidae) iz severne tunizijske obale (osrednje Sredozemsko morje)*

FLORA  
 FLORA  
 FLORA

**Martina ORLANDO-BONACA, Erik LIPEJ, Romina BONACA & Leon Lojze ZAMUDA**

- Improvement of the Ecological Status of the *Cymodocea nodosa* Meadow near the Port of Koper ..... 185  
*Izboljšanje ekološkega stanja morskega travnika kolenčaste cimodoceje (*Cymodocea nodosa*) v bližini koprskega pristanišča*

FAVNA  
 FAVNA  
 FAVNA

**Manja ROGELJA, Martin VODOPIVEC & Alenka MALEJ**

- Cestum veneris* Lesueur, 1813 (Ctenophora) – a Rare Guest in the Northern Adriatic Sea ..... 197  
*Cestum veneris* Lesueur, 1813 (Ctenophora) – redenk gost v severnem Jadranu

**Adla KAHRić, Dalila DELIĆ & Dejan KULIJER**

- Notospermus annulatus* (Nemertea: Lineidae), a New Record for Bosnia and Herzegovina ..... 205  
*Notospermus annulatus* (Nemertea: Lineidae), prvi zapis o pojavljanju za Bosno in Hercegovino

**Andrea LOMBARDO & Giuliana MARLETTA**

- Report of an Interesting *Trapania* (Gastropoda: Nudibranchia: Goniodorididae) Specimen from Central Eastern Sicily ..... 211  
*Zapis o zanimivem primerku iz rodu Trapania (Gastropoda: Nudibranchia: Goniodorididae) iz osrednje vzhodne Sicilije*

**Abdelkarim DERBALI & Othman JARBOUI**

- Stock Assessment, Cartography and Sexuality of the Wedge Clam *Donax trunculus* in the Gulf of Gabes (Tunisia) ..... 217  
*Ocena staleža, kartografija in spolnost klinaste školjke Donax trunculus v gabeškem zalivu (Tunizija)*

**Abdelkarim DERBALI, Aymen HADJ TAIEB & Othman JARBOUI**

- Length-Weight Relationships and Density of Bivalve Species in the Shellfish Production Area of Zarzis (Tunisia, Central Mediterranean Sea) ..... 229  
*Dolžinsko-masni odnos in gostota školjk na gojišču školjk v predelu Zarsisa (Tunizija, osrednje Sredozemsko morje)*

**Toni KOREN**

- The Diversity of Moths (Lepidoptera: Heterocera) of Significant Landscape Donji Kamenjak and Medulin Archipelago, Istria, Croatia ..... 237  
*Raznolikost nočnih metuljev (Lepidoptera: Heterocera) Pomembne pokrajine Donji Kamenjak in Medulinski arhipelag, Istra, Hrvaška*

OCENE IN PEROČILA  
 RECENSIONI E RELAZIONI  
 REVIEWS AND REPORTS

**Ines Mandić Mulec & Nives Ogrinc**

- Recenzija knjige: Mikrobična biogeokemijska voda ..... 263  
 Kazalo k slikam na ovtiku ..... 265  
*Index to images on the cover* ..... 265



received: 2022-04-11

DOI 10.19233/ASHN.2022.04

## MORPHOMETRIC MEASUREMENTS OF THE YOUNG OF THREE CARCHARHINID SPECIES FROM NORTHEASTERN LEVANT (MEDITERRANEAN SEA)

*Deniz ERGÜDEN*

Marine Sciences Department, Marine Sciences and Technology Faculty, İskenderun Technical University, İskenderun, Hatay, Turkey

*Deniz AYAS*

Faculty of Fisheries, Mersin, Turkey

*Hakan KABASAKAL*

Ichthyological Research Society, Ümraniye, İstanbul, Turkey

e-mail: kabasakal.hakan@gmail.com

### ABSTRACT

*This study presents the morphometric measurements of young individuals of Carcharhinus altimus (Springer, 1950), C. obscurus (Lesueur, 1818), and C. plumbeus (Nardo, 1827). These data provide valuable information on the morphometry of the populations of the three species in the northeastern Levant.*

**Key words:** *Carcharhinus, morphometry, Levantine Sea, Turkey*

## MISURE MORFOMETRICHE DEI GIOVANI DI TRE SPECIE DI CARCARINIDI DEL LEVANTE NORD-ORIENTALE (MAR MEDITERRANEO)

### SINTESI

*Lo studio presenta le misure morfometriche di giovani individui di Carcharhinus altimus (Springer, 1950), C. obscurus (Lesueur, 1818) e C. plumbeus (Nardo, 1827). Questi dati forniscono informazioni preziose sulla morfometria delle popolazioni delle tre specie nel Levante nord-orientale.*

**Parole chiave:** *Carcharhinus, morfometria, Mar Levantino, Turchia*

## INTRODUCTION

Since the genus *Carcharhinus* includes some of the largest species of sharks in the Mediterranean waters, a remarkable research effort has been spent by many authors to provide new data for a better understanding of the species of this genus in the entire region (e.g., Capapé, 1984; Hemida et al., 2002; Lipej et al., 2008; Başusta, 2016; Filiz, 2018; Capapé et al., 2018; Azab et al., 2019). Due to their vital role in describing fish species (Moyle & Cech, Jr., 1988), morphometric measurements have been

reported in most of these studies, and therefore, a remarkable archive of Mediterranean *Carcharhinus* spp. morphometrics is available.

In the present study, the authors report the morphometrics of three species belonging to the genus *Carcharhinus* Blainville, 1816; the bignose shark, *Carcharhinus altimus* (Springer, 1950); the dusky shark, *C. obscurus* (Lesueur, 1818); and the sandbar shark, *C. plumbeus* (Nardo, 1827), which were incidentally captured in the coastal waters of northeastern Levant. The authors also aim to contribute to the available data-set of morphometric measurements



**Fig. 1:** Map depicting the capture localities of the examined *Carcharhinus altimus* (●), *C. obscurus* (▲) and *C. plumbeus* (■) specimens. Square in the small map indicates the locality in the eastern Levant covered by the present study.  
**Sl. 1:** Zemljevid obravnavanega območja z označenimi lokalitetami, kjer so bili ujeti primerki vrst *Carcharhinus altimus* (●), *C. obscurus* (▲) in *C. plumbeus* (■). Kvadrat na manjšem zemljevidu označuje območje vzhodnega Levanta, obravnavanega v pričujoči raziskavi.

with records of these three Mediterranean carcharhinid species, which may be useful for future research of the same objective.

## MATERIAL AND METHODS

Young individuals of *Carcharhinus altimus* (Springer, 1950) (n=6; TL range 54.5–113 cm, mean  $73.1 \pm 22.3$  cm), *C. obscurus* (Lesueur, 1818) (n=1; TL 64.5 cm), and *C. plumbeus* (Nardo, 1827) (n=1; TL 91 cm), were incidentally captured by coastal artisanal long-liners, trammel-netters and gill-netters along the coastline stretching between Dana Island and the mouth of the Göksu River (Fig. 1). *C. plumbeus* is a species protected by the current fisheries law of Turkey and any living sandbar sharks accidentally captured should be immediately released back into the sea. Although not included in the list of protected marine species, based on the researchers' decision, all alive individuals of *C. altimus* and *C. obscurus* in this study were also released due to their conservation status (Data Deficient; Serena et al., 2020). Only dead individuals were retained on board and landed. Before performing any observations and incisions on the sharks, the ethic committee approval was acquired for each shark, which was stored in the freezer.

Species identification follows Grace (2001) and Serena (2005), and taxonomic nomenclature follows Serena et al. (2020). Measurements were recorded on thawed individuals. Total length (TL) is the distance between the tip of the snout and the tip of the upper caudal lobe, where the upper lobe is in stretched position (Ebert & Stehmann, 2013). Twenty-seven morphometric measurements were taken using a measurement tape and recorded to the nearest 0.1 mm, following the procedure outlined in Ebert and Stehmann (2013). Previously published morphometric measurements of the same species were extracted from the relevant references (Devaraj & Gulati, 2004; Lipej et al., 2008; Capapé et al., 2018; Azab et al., 2019; Lee et al., 2019; Ayas et al., 2020; Ergüden et al., 2020; Turan et al., 2020; Froese & Pauly, 2022). All of the measurements were presented as percentages of TL, and for species with multiple individuals, percentages of TL values were computed based on the mean ( $\pm$ SD) of each measurement. The captured individuals were recorded in the Museum of Marine Life, Mersin University with catalogue numbers (MEUFC-22-11-135 - MEUFC-22-11-142). Excel worksheets of the three species are available from the corresponding author for inspection on request.

## RESULTS AND DISCUSSION

The main morphological characters allowing us to distinguish the three species of *Carcharhinus* were as follows (Ebert & Stehmann, 2013):

### *Carcharhinus altimus* (Springer, 1950)

Anterior nasal flaps usually high and triangular. Distance from nostrils to mouth less than 2.4 times the mouth width. Upper anterolateral teeth very high; upper anterolateral teeth usually in 15 rows. First dorsal fin lower, with height much less than half predorsal space. Interdorsal ridge high.

### *Carcharhinus obscurus* (Lesueur, 1818)

First dorsal fin origin over or anterior to pectoral fin free rear tips. Pectoral fins large and falcate. Serrations of upper anterolateral teeth small and not very coarse. Inner margin of second dorsal fin shorter and generally less than twice fin height (up to 2.1 times). Interdorsal ridge low.

### *Carcharhinus plumbeus* (Nardo, 1827)

Anterior nasal flaps usually low and inconspicuous. Distance from nostrils to mouth more than 2.4 times the mouth width. Upper anterolateral teeth moderately high; upper anterolateral teeth usually in 14 rows. First dorsal fin very high, with height about half predorsal space. Interdorsal ridge low.

In a recent review of the distribution of chondrichthyans in the Mediterranean Sea, Serena et al. (2020) described *C. altimus* as a vagrant species for the Mediterranean Sea, and *C. obscurus* and *C. plumbeus* as resident carcharhinids in the region. The occurrence of these three species in Turkish waters has been mostly reported from Mediterranean coasts, while the distribution of *C. plumbeus* in the region also extends to southern Aegean waters (Kabasakal, 2021).

Since all of the individuals examined had prominent interdorsal ridges (Fig. 2), the *C. brachyurus*, *C. brevipinna*, *C. limbatus*, and *C. melanopterus* species were easily excluded as possibilities. Unlike *C. falciformis*, in which the free rear tip of the 2<sup>nd</sup> dorsal fin is very long ( $\geq 2$  to 3 times in 2<sup>nd</sup> D fin height), the length of the mentioned body part in the examined individuals was shorter (Tables 1–3); as a result, *C. falciformis*, although also exhibiting an interdorsal ridge, was also excluded as a possibility. Six out of the eight examined sharks coincided with the descriptions of *C. altimus*, and the remaining two individuals were identified as *C. obscurus* and *C. plumbeus*, respectively (Grace, 2001; Serena, 2005).

Despite the slight differences, our measurements generally agree with those given by Compagno (1984) for *Carcharhinus* spp. occurring worldwide, and Ebert & Stehmann (2013) for species occurring in the North Atlantic. The percentage of eye length in TL (EYL-TL) in *C. obscurus* (1.8%) fell within the range given by Compagno (1984) and Ebert & Stehmann (2013), which is reported as 1–2.1%. The EYL-TL ratio of *C. altimus* for the mean and maximum



**Fig. 2:** Some of the specimens of *Carcharhinus* spp. examined in the present study: (a) ventral view of the head and whole lateral view of *C. altimus*; (b) dorsal view of the head and whole lateral view of *C. obscurus*; and (c) ventral view of the head and whole lateral view of *C. plumbeus*. Scale bar = 10 cm.

**Sl. 2:** Nekaj preiskanih primerkov iz rodu *Carcharhinus* iz te raziskave: a) pogled na glavo vrste *C. altimus* s trebušne strani in pogled z boka; (b) pogled na glavo vrste *C. obscurus* s hrbitne strani in pogled z boka; in (c) pogled na glavo vrste *C. plumbeus* s trebušne strani in pogled z boka. Merilo = 10 cm.

values of the examined specimens (1.5% and 1.7%, respectively), were in agreement with those reported by Compagno (1984) and Ebert & Stehmann (2013), which is 1.4–2.3%; however, the percentage of eye length in TL in the smallest examined specimen (1.2%) was lower than the reported range. In the examined specimen of *C. plumbeus*, the percentage of eye length in TL (1.2%) was also lower than the reported range (1.7–2.9%; Compagno, 1984; Ebert & Stehmann, 2013).

Compagno (1984) and Ebert & Stehmann (2013) reported the percentages of 2<sup>nd</sup> dorsal fin height in TL (D2H-TL) and pectoral anterior margin length in

TL (PAL-TL) in *Carcharhinus altimus* as 2.8–3.4% and 20–22%, respectively. Mean and maximum values of D2H-TL and PAL-TL ratios of the examined specimens of *C. altimus* (2.9% and 3.1% for mean and max. D2H-TL, and 17.7% and 17.8% for mean and max. PAL-TL, respectively), coincided with the reported percentages. On the other hand, the same percentages of the smallest examined specimen (2.6% for D2H-TL and 16.5% for PAL-TL) are lower than the above-published percentage ranges. D2H-TL percentages for examined specimens of *C. obscurus* and *C. plumbeus* (2.3% and 3.3%, respectively), are agreed with those reported by Compagno (1984) and Ebert & Stehmann (2013),

**Tab. 1: Morphometric measurements of *Carcharhinus altimus* examined in the present study. In bold the percentages mentioned in Compagno (1984) and Ebert & Stehmann (2013).****Tab. 1: Morfometrične meritve na primerkih vrste *Carcharhinus altimus*, preiskanih v pričujoči raziskavi. V mstnem tisku so deleži, ki jih omenjajo Compagno (1984) in Ebert & Stehmann (2013).**

| Measurements                | Present Study<br>(TL range 54.5–113 cm) |      |                    |                   |                   | Turan et al.<br>(2020) |             | Ayas et al.<br>(2020) |             | Azab et al.<br>(2019) |            | Froese<br>& Pauly<br>(2022)<br>(74 cm TL) |
|-----------------------------|---|------|--------------------|-------------------|-------------------|------------------------|-------------|-----------------------|-------------|-----------------------|------------|---|
|                             | Mean                                    | SD   | % of<br>mean<br>TL | % of<br>min<br>TL | % of<br>max<br>TL | TL                     | % of<br>TL  | TL                    | % of<br>TL  | TL                    | % of<br>TL | % of TL                                   |
| Total length (TL)           | 73.1                                    | 22.3 |                    |                   |                   | 68                     |             | 65.2                  |             | 54.6                  |            |   |
| Fork length (FL)            | 58.4                                    | 18.0 | 79.9               | 79.8              | 80.5              |                        |             | 53.0                  | 81.3        |                       | 78.7       | 82.5                                      |
| Standard length             | 53.3                                    | 16.5 | 72.9               | 72.5              | 73.5              | 59.0                   | 86.8        | 48.5                  | 74.4        |                       | 70.7       | 79.6                                      |
| Head length                 | 18.1                                    | 4.8  | 24.7               | 26.4              | 23.9              | 11.1                   | 16.3        | 15.8                  | 24.2        |                       | 17.5       |   |
| Mouth length                | 7.0                                     | 2.1  | 9.6                | 10.1              | 9.3               |                        |             | 5.6                   | 8.6         |                       |            |   |
| Eye length                  | 1.1                                     | 0.2  | <b>1.5</b>         | <b>1.7</b>        | <b>1.2</b>        | 0.8                    | <b>1.2</b>  | 0.9                   | <b>1.4</b>  |                       |            |   |
| Eye height                  | 1.1                                     | 0.2  | 1.4                | 1.7               | 1.2               |                        |             |                       |             |                       |            |   |
| Internarial<br>distance     | 4.5                                     | 1.1  | 6.1                | 7.3               | 5.7               |                        |             |                       |             |                       |            |   |
| Pre-branchial<br>length     | 14.8                                    | 4.1  | 20.3               | 20.2              | 19.5              | 9.0                    | 13.2        | 13.5                  | 20.7        |                       |            | 23.5                                      |
| Pre-orbital length          | 6.8                                     | 1.9  | 9.3                | 8.8               | 8.8               |                        |             | 5.8                   | 8.9         |                       |            |   |
| Pre-D1 fin length           | 22.8                                    | 6.8  | 31.1               | 31.2              | 31.0              |                        |             | 21.0                  | 32.2        |                       |            | 29.9                                      |
| Pre-D2 fin length           | 46.9                                    | 14.3 | 64.1               | 62.4              | 63.7              |                        |             |                       |             |                       |            |   |
| Pre-pectoral fin<br>length  | 17.8                                    | 4.4  | 24.4               | 24.8              | 22.1              |                        |             | 13.5                  | 20.7        |                       |            | 24  |
| D1 fin length               | 11.6                                    | 4.1  | 15.8               | 15.4              | 16.8              |                        |             | 9.6                   | 14.7        |                       |            |   |
| D1 fin height               | 7.9                                     | 2.9  | 10.8               | 9.2               | 11.1              |                        |             | 6.4                   | 9.8         |                       |            |   |
| D1 fin base                 | 8.3                                     | 2.8  | 11.4               | 11.0              | 11.9              |                        |             | 7.0                   | 10.7        |                       |            |   |
| D2 fin length               | 5.1                                     | 1.2  | 6.9                | 7.0               | 4.9               |                        |             | 4.5                   | 6.9         |                       |            |   |
| D2 fin height               | 2.2                                     | 0.9  | <b>2.9</b>         | <b>2.6</b>        | <b>3.1</b>        | 2.0                    | <b>2.9</b>  |                       |             |                       |            |   |
| D2 fin base                 | 2.9                                     | 1.1  | 3.9                | 4.4               | 4.0               |                        |             | 2.1                   | 3.2         |                       |            |   |
| Pectoral fin<br>length      | 13.0                                    | 4.3  | <b>17.7</b>        | <b>16.5</b>       | <b>17.8</b>       | 10.5                   | <b>15.4</b> | 13.5                  | <b>20.7</b> |                       |            |   |
| Pectoral fin base           | 5.3                                     | 2.4  | 7.3                | 6.4               | 8.8               |                        |             | 2.8                   | 4.3         |                       |            |   |
| Pelvic fin length           | 5.8                                     | 2.0  | 8.0                | 7.3               | 8.0               |                        |             |                       |             |                       |            |   |
| Pelvic fin base             | 2.6                                     | 0.9  | 3.6                | 3.3               | 3.5               |                        |             | 2.8                   | 4.3         |                       |            |   |
| Anal fin length             | 5.8                                     | 2.0  | 7.9                | 8.3               | 8.0               |                        |             |                       |             |                       |            |   |
| Anal fin base               | 3.4                                     | 1.2  | 4.6                | 4.6               | 4.9               |                        |             | 2.4                   | 3.7         |                       |            |   |
| Caudal upper<br>lobe length | 19.4                                    | 6.2  | 26.6               | 26.6              | 27.0              |                        |             | 16.0                  | 24.5        |                       |            |   |
| Caudal lower<br>lobe length | 8.1                                     | 2.6  | 11.1               | 12.8              | 11.1              |                        |             | 6.2                   | 9.5         |                       |            |   |

**Tab. 2: Morphometric measurements of *Carcharhinus obscurus* examined in the present study. In bold the percentages mentioned in Compagno (1984) and Ebert & Stehmann (2013). TL of respective specimens examined by Devaraj & Gulati (2004), Azab et al. (2019) and Lee et al. (2019) are given in parentheses under the authors' names.**

**Tab. 2: Morfometrične meritve na primerku vrste *Carcharhinus obscurus*, preiskanem v pričujoči raziskavi. V mestnem tisku so deleži, ki jih omenjajo Compagno (1984) in Ebert & Stehmann (2013). Dolžine telesa primerkov, ki jih navajajo Devaraj & Gulati (2004), Azab et al. (2019) in Lee in sod. (2019) so podane v oklepajih pod avtorjevimi imeni.**

| Measurements             | Present Study |             | Devaraj & Gulati<br>(2004)<br>(99.5 cm TL) | Azab et al. (2019)<br>(58.08 cm TL) | Lee et al. (2019)<br>(83.9 cm TL) |
|--------------------------|---------------|-------------|--|-------------------------------------|-----------------------------------|
|                          | cm            | % of TL     | % of TL                                    | % of TL                             | % of TL                           |
| Total length (TL)        | 64.5          |             |  |                                     |                                   |
| Fork length (FL)         | 51            | 79.1        |  |                                     | 78.6                              |
| Standard length          | 47.5          | 73.6        | 73.2                                       | 73.8                                | 71.2                              |
| Head length              | 16.5          | 25.6        | 18.5                                       |                                     | 18.8                              |
| Mouth length             | 6.5           | 10.1        | 7.8  |                                     |                                   |
| Eye length               | 1.15          | <b>1.8</b>  | <b>1.7</b>                                 | <b>1.7</b>                          |                                   |
| Eye height               | 1.05          | 1.6         |  |                                     |                                   |
| Internarial distance     | 3.9           | 6.0         |  |                                     |                                   |
| Pre-branchial length     | 13            | 20.2        | 18.5                                       | 5.4                                 |                                   |
| Pre-orbital length       | 6             | 9.3         | 8.1  |                                     |                                   |
| Pre-D1 fin length        | 19            | 29.5        | 31.0                                       | 34.2                                |                                   |
| Pre-D2 fin length        | 41            | 63.6        | 62.1                                       | 63.9                                |                                   |
| Pre-pectoral fin length  | 16            | 24.8        | 21.2                                       | 22.6                                |                                   |
| D1 fin length            | 10            | 15.5        |  |                                     |                                   |
| D1 fin height            | 6.5           | 10.1        | 8.1  | 11.2                                |                                   |
| D1 fin base              | 8             | 12.4        | 8.6  | 82.4                                |                                   |
| D2 fin length            | 4.5           | 7.0         |  | 3.0                                 |                                   |
| D2 fin height            | 1.5           | <b>2.3</b>  | <b>2.0</b>                                 |                                     |                                   |
| D2 fin base              | 2.5           | 3.9         | 2.8  | 2.2                                 |                                   |
| Pectoral fin length      | 10            | <b>15.5</b> | <b>11.5</b>                                | <b>16.1</b>                         |                                   |
| Pectoral fin base        | 5             | 7.8         | 5.2  | 5.9                                 |                                   |
| Pelvic fin length        | 5             | 7.8         |  | 4.0                                 |                                   |
| Pelvic fin base          | 2             | 3.1         | 4.2  |                                     |                                   |
| Anal fin length          | 4.9           | 7.6         |  | 4.7                                 |                                   |
| Anal fin base            | 2.5           | 3.9         | 3.6  | 2.8                                 |                                   |
| Caudal upper lobe length | 17            | 26.4        | 25.5                                       | 27.1                                |                                   |
| Caudal lower lobe length | 7             | 10.9        | 10.6                                       | 11.8                                |                                   |

**Tab. 3: Morphometric measurements of *Carcharhinus plumbeus* examined in the present study. In bold the percentages mentioned in Compagno (1984) and Ebert & Stehmann (2013).****Tab. 3: Morfometrične meritve na primerku vrste *Carcharhinus plumbeus*, opravljene v pričujoči raziskavi. V mastnem tisku so deleži, ki jih omenjajo Compagno (1984) in Ebert & Stehmann (2013).**

| Measurements             | Present Study |            | Lipej et al. (2008) |     |              | Capapé et al. (2018) |             | Ergüden et al. (2020) |             |
|--------------------------|---------------|------------|---------------------|-----|--------------|----------------------|-------------|-----------------------|-------------|
|                          | cm            | % of TL    | Mean                | SD  | % of mean TL | cm                   | % of TL     | cm                    | % of TL     |
| Total length (TL)        | 91            |            | 74.3                | 6.2 |              | 89                   |             | 68                    |             |
| Fork length (FL)         | 72.5          | 79.7       | 59.8                | 5.4 | 80.5         |                      |             |                       |             |
| Standard length          | 65            | 71.4       | 54.7                | 5.1 | 73.6         |                      |             |                       |             |
| Head length              | 22.5          | 24.7       | 19.1                | 1.9 | 25.7         | 17                   | 19.1        | 16.7                  | 24.6        |
| Mouth length             | 8             | 8.8        | 7.2                 | 0.5 | 9.6          | 8                    | 9.0         |                       |             |
| Eye length               | 1.1           | <b>1.2</b> | 1.0                 | 0.2 | <b>1.4</b>   | 1.1                  | <b>1.2</b>  | 0.9                   | <b>1.3</b>  |
| Eye height               | 0.9           | 1.0        | 1.1                 | 0.1 | 1.4          | 1.2                  | 1.3         |                       |             |
| Internarial distance     | 5.3           | 5.8        | 4.5                 | 0.3 | 6.0          | 5                    | 5.6         |                       |             |
| Pre-branchial length     | 19            | 20.9       | 14.8                | 1.6 | 19.9         | 16.5                 | 18.5        |                       |             |
| Pre-orbital length       | 8.5           | 9.3        | 6.5                 | 1.0 | 8.7          |                      |             |                       |             |
| Pre-D1 fin length        | 28.5          | 31.3       | 21.5                | 1.7 | 28.9         |                      |             |                       |             |
| Pre-D2 fin length        | 59            | 64.8       | 46.7                | 3.8 | 62.8         |                      |             |                       |             |
| Pre-pectoral fin length  | 23            | 25.3       | 17.2                | 2.0 | 23.1         |                      |             |                       |             |
| D1 fin length            | 14.5          | 15.9       | 10.4                | 0.3 | 14.0         | 12                   | 13.5        | 12                    | 17.6        |
| D1 fin height            | 10.5          | 11.5       | 7.2                 | 1.0 | 9.6          |                      |             |                       |             |
| D1 fin base              | 10            | 11.0       | 9.0                 | 0.3 | 12.1         | 10                   | 11.2        |                       |             |
| D2 fin length            | 7             | 7.7        | 3.1                 | 0.3 | 4.2          | 4                    | 4.5         | 5.4                   | 7.9         |
| D2 fin height            | 3             | <b>3.3</b> | 2.3                 | 0.5 | <b>3.1</b>   |                      |             |                       |             |
| D2 fin base              | 3.5           | 3.8        | 2.9                 |     | 3.9          | 4                    | 4.5         |                       |             |
| Pectoral fin length      | 15.5          | <b>17</b>  | 12.5                | 0.7 | <b>16.9</b>  | 14                   | <b>15.7</b> | 9.2                   | <b>13.5</b> |
| Pectoral fin base        | 7             | 7.7        | 4.9                 | 0.6 | 6.6          | 5.5                  | 6.2         |                       |             |
| Pelvic fin length        | 7             | 7.7        | 6.0                 | 0.6 | 8.0          | 4                    | 4.5         | 5.8                   | 8.5         |
| Pelvic fin base          | 3.5           | 3.8        | 3.1                 | 0.3 | 4.1          | 4                    | 4.5         |                       |             |
| Anal fin length          | 7             | 7.7        | 5.6                 | 0.3 | 7.6          | 5                    | 5.6         | 5.3                   | 7.8         |
| Anal fin base            | 4             | 4.4        | 3.1                 | 0.3 | 4.2          | 3.5                  | 3.9         |                       |             |
| Caudal upper lobe length | 25            | 27.5       | 10.6                | 1.1 | 14.3         | 22                   | 24.7        |                       |             |
| Caudal lower lobe length | 9             | 9.9        | 5.2                 | 0.5 | 7.0          | 8                    | 9.0         | 6                     | 8.8         |

of which as 1.5–2.3% for *C. obscurus* and 2.1–3.5% for *C. plumbeus*. Similarly, PAL-TL percentages of *C. obscurus* and *C. plumbeus* (1.5% and 17%, respectively) were within the ranges reported for *C. obscurus* (17–22%) and *C. plumbeus* (17–22%) by Compagno (1984) and Ebert & Stehmann (2013).

The slight variations between our measurements and those given by Compagno (1984) and Ebert & Stehmann (2013) suggest incongruity with the published descriptive ratios; however, similar variations have been observed in young individuals of the present *Carcharhinus* spp. before. When comparing our results to those reported for young individuals of *C. altimus*, *C. obscurus* and *C. plumbeus* from several regions of the Mediterranean Sea and elsewhere in the world (Devaraj & Gulati, 2004; Lipej et al., 2008; Capapé et al., 2018; Azab et al., 2019; Lee et al., 2019; Ayas et al., 2020; Ergüden et al., 2020; Turan et al., 2020; Froese & Pauly, 2022; Tables 1–3) it is clear that the results concerning the three examined carcharhinid species from our study coincide better with the percentages reported by the above authors for young specimens (bold numbers in the relevant tables) than they do with those reported by Compagno (1984) and Ebert & Stehmann (2013) for adult individuals.

The percentage of EYL-TL of the examined specimens of *Carcharhinus altimus* was in agreement with those reported by Ayas et al. (2020) and Turan et al. (2020) (Table 1). The percentage range of D2H-TL of the examined specimens of *C. altimus* was also within the range reported by Turan et al. (2020); however, the percentage of PAL-TL of the examined specimens exceeded the percentage range reported by Ayas et al. (2020), and was higher than the ratio reported by Turan et al. (2020) (Table 1). Specimens of *C. altimus* reported by Ayas et al. (2020) and Turan et al. (2020) were also incidentally captured in the vicinity of the present sampling localities.

The EYL-TL percentages in the present specimen of *Carcharhinus obscurus* and in the Indian Ocean specimen reported by Devaraj & Gulati (2004) are in agreement with the percentage range reported for dusky sharks from Korean waters (Lee et al., 2019) (Table 2), and the percentage of D2H-TL for the present specimen fits within the range reported by Lee et al. (2019), as well (Table 2). The PAL-TL percentages reported for the present specimen (15.5%) and for the Indian Ocean specimen (16.1%; Devaraj & Gulati, 2004) both exceed the range recorded in Korean dusky sharks (11.5–13.9%; Lee et al., 2019).

For *Carcharhinus plumbeus*, the EYL-TL, D2H-TL, and PAL-TL percentages recorded in the present speci-

men differ from those recorded in specimens reported by Lipej et al. (2008), Capapé et al. (2018) and Ergüden et al. (2020), standing at 1.2 vs. 1.4%, 3.1 vs. 3.3% and 13.5 vs. 17%, respectively (Table 3).

## CONCLUSIONS

Due to similar body shapes, colours and overlapping distributions, the Carcharhinidae genera can generally be difficult to identify, particularly *Carcharhinus* spp. (Grace, 2001). Since the existing literature addressing worldwide distribution (e.g., Compagno, 1984) or covering wide marine regions (e.g., Serena, 2005; Ebert & Stehmann, 2013) may not be applicable to carcharhinids occurring in Turkish waters, a set of multiple descriptions including a broad range of ratios and characters may be needed for their identification, as was the case in the present study. Identification of *Carcharhinus* spp. is facilitated by division into two groups based on the presence or absence of an interdorsal ridge. The origin of the first dorsal fin in relation to the pectoral fin and also the snout shape are useful for further subdivision of each group and, ultimately, species identification. This conventional order of division of characteristics follows the accepted identification keys (e.g., Compagno, 1984; Grace, 2001; Serena, 2005; Ebert & Stehmann, 2013), so what could be the underlying reasons for the differences in morphometric percentages?

Geographically distant (allopatric) populations of the same fish species tend to exhibit morphometric characters at the opposite margins of the value ranges (Cailliet et al., 1986). The morphometric differences seen in the present study can thus be considered admissible based on the above-mentioned situation. Although the morphometric data reported for *Carcharhinus altimus*, *C. obscurus* and *C. plumbeus* in the present study provide valuable information on the morphology of their populations in the northeastern Levant, it is clear that morphometric analysis of more specimens of each *Carcharhinus* species are required to obtain statistically significant results.

## ACKNOWLEDGMENTS

The authors thank to the coastal artisanal fishermen for donating the incidentally captured specimens of *Carcharhinus* spp. and releasing the alive sharks as fast as possible. Very special thanks go to two anonymous referees for their valuable comments, which improved the content and quality of the article.

## MORFOMETRIČNE MERITVE MLADIČEV TREH VRST MORSKIH PSOV IZ DRUŽINE CARCHARHINIDAE IZ SEVERNOVZHODNEGA LEVANTA (SREDOZEMSKO MORJE)

Deniz ERGÜDEN

Marine Sciences Department, Marine Sciences and Technology Faculty, İskenderun Technical University, İskenderun, Hatay, Turkey

Deniz AYAS

Faculty of Fisheries, Mersin, Turkey

Hakan KABASAKAL

Ichthyological Research Society, Ümraniye, İstanbul, Turkey

e-mail: kabasakal.hakan@gmail.com

### POVZETEK

Avtorji poročajo o morfometričnih meritvah na mladih primerkih vrst *Carcharhinus altimus* (Springer, 1950), *C. obscurus* (Lesueur, 1818) in *C. plumbeus* (Nardo, 1827). Te meritve nudijo pomembne podatke o morfometriji populacij treh vrst v severovzhodnem Levantu.

**Ključne besede:** *Carcharhinus*, morfometrija, Levantsko morje, Turčija

## REFERENCES

- Ayas, D., N. Çiftçi, E. Yalçın, H. D. Akbora, M. Bakan & D. Ergüden (2020):** First record of the big nose shark, *Carcharhinus altimus* (Springer, 1950) from Mersin bay. IJFAS, 8, 132-136.
- Azab, A.M., H.M.M. Khalaf-Allah, M.M.H. Sarhan & M.A.M. El-Tabakh (2019):** Carcharhinid shark species (Family Carcharhinidae), with special reference to the first records in the Egyptian Mediterranean waters, Alexandria, Egypt. Egypt. J. Aquat. Biol. Fish., 23, 545-559.
- Başusta, N. (2016):** Length-weight relationship of sandbar shark *Carcharhinus plumbeus* (Nardo, 1827) in İskenderun Bay (north-eastern Mediterranean Sea). Rapp. Comm. int. Mer Médit., 41, 318.
- Cailliet, G.M., M.S. Love & A.W. Ebeling (1986):** Fishes. A Field and Laboratory Manual on Their Structure, Identification, and Natural History. Wadsworth Publishing Company, Belmont, 194 pp.
- Capapé, C. (1984):** Nouvelles données sur la morphologie et la biologie de la reproduction de *Carcharhinus plumbeus* (Nardo, 1827) (Pisces, Carcharhinidae) des côtes Tunisiennes. Inv. Pesq., 48, 115-137.
- Capapé, C., S. Rafrafi-Nouria, K. Ounifi-Ben Amor & M.M. Ben Amor (2018):** Additional records of sandbar shark, *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae) from the northern Tunisian coast (central Mediterranean Sea). Annales, Ser. Hist. Nat., 28, 99-104.
- Compagno, L.J.V. (1984):** FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2 - Carcharhiniformes. FAO Fish. Synop. 125(4/2), FAO, Rome, pp. 251-655.
- Devaraj, M. & D. Gulati (2004):** Morphometry and biology of the dusky shark, *Carcharhinus obscurus* (LeSueur), from the oceanic region off the west coast of India. In: V.S. Somvanshi (ed): Large Marine Ecosystems: Exploration and Exploitation for Sustainable Development and Conservation on Fish Stocks. Fishery Survey of India, Ministry of Agriculture, Mumbai, pp. 100-112.
- Ebert, D.A. & M.F.W. Stehmann (2013):** Sharks, batoids, and chimaeras of the North Atlantic FAO Species Catalogue for Fishery Purposes. No. 7. Rome, FAO, 523 pp.
- Ergüden, D., H. Kabasakal & F. Kabaklı (2020):** Young-of-the-year sandbar shark, *Carcharhinus plumbeus* (Nardo, 1827) (Carcharhiniformes: Carcharhinidae), caught in İskenderun Bay. FishTaxa, 18, 18-22.
- Filiz, H. (2018):** Year-round aggregation of sandbar shark, *Carcharhinus plumbeus* (Nardo, 1827), in Boncuk Cove in the southern Aegean Sea, Turkey (Carcharhiniformes: Carcharhinidae). Zool. Middle East, 65, 35-39.
- Froese, R. & D. Pauly (Eds.) (2022):** FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org). version (02/2022) (Last accession: 7 April 2022).
- Grace, M. (2001):** Field guide to requiem sharks (Elasmobranchimorphi: Carcharhinidae) of the western north Atlantic. U. S. dep. Commer., NOAA Tech. Rep. NMFS 153, 32 pp.
- Hemida, F., R. Seridji, N. Labidi, J. Bensaci & C. Capapé (2002):** Records of *Carcharhinus* spp. (Chondrichthyes: Carcharhinidae) from off the Algerian coast (southern Mediterranean). Acta Adriat., 43, 83-92.
- Kabasakal, H. (2021):** A review of shark biodiversity in Turkish waters. updated inventory, new arrivals, questionable species, and conservation issues. Annales, Ser. Hist. Nat., 31, 181-194.
- Lee, W.J., Y. Kim, J.-H. Na & J.-K. Kim (2019):** First reliable record of the dusky shark *Carcharhinus obscurus* (Lesueur, 1818) (Carcharhinidae, Pisces) from Korea. KFAS 52, 696-701.
- Lipej, L., B. Mavrič, Ž. Dobrajc & C. Capapé (2008):** On the occurrence of the sandbar shark, *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae) off the Slovenian coast (northern Adriatic). Acta Adriat., 49, 137-145.
- Moyle, P.B. & J.J. Cech, Jr. (1988):** Fishes. An Introduction to Ichthyology. Prentice Hall, Englewood Cliffs, 559 pp.
- Serena, F. (2005):** Field identification guide to the sharks and rays of the Mediterranean and Black Sea. FAO Species Identification Guide for Fishery Purposes. Rome, FAO, 97 pp.
- Serena, F., A.J. Abella, F. Bargnesi, M. Barone, F. Colloca, F. Ferretti, F. Fiorentino, J. Jenrette & S. Moro (2020):** Species diversity, taxonomy and distribution of Chondrichthyes in the Mediterranean and Black Sea. Eur. Zool. J., 87, 497-536, DOI: 10.1080/24750263.2020.1805518.
- Turan, C., M. İğde, A. Ergenler & D. Ayas (2020):** New record of the bignose shark, *Carcharhinus altimus* (Springer, 1950) (Carcharhiniformes: Carcharhinidae), in Turkish waters. J. Black Sea/Medit., 26, 119-125.