

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnsič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Založnika/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 6. 12. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija *Annales, Series Historia Naturalis* je vključena v naslednje podatkovne baze / La rivista *Annales, series Historia Naturalis* è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / CONTENTSSREDOZEMSKÉ HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Hakan KABASAKAL**

Analysis of Confirmed Shark Attacks
in the Eastern Mediterranean Sea and
the Sea of Marmara (1827–2025) 169
Analiza potrjenih napadov morskih psov
in Marmarskem morju (1827–2025)

Alen SOLDO

Observations of a Juvenile Basking Shark
Cetorhinus maximus in the Adriatic Sea
Support the Hypothesis of a Distinct
Mediterranean Population 187
Opazovanja mladege morskega psa
orjaka (Cetorhinus maximus)
v Jadranskem morju podpirajo
hipotezo o ločeni sredozemski
populaciji

**Jacopo BERNARDI, Pero UGARKOVIĆ &
Ilija ČETKOVIĆ**

An Unusual Encounter with a Juvenile Kitefin
Shark, *Dalatias licha*, in Shallow Coastal
Waters of Montenegro (Adriatic Sea) 197
Nenavadno srečanje z mladim primerkom
klinoplavutega morskega psa, Dalatias licha,
v plitvih obalnih vodah Črne gore
(Jadransko morje)

**Cemal TURAN, Alen SOLDO, Servet A. DOĞDU,
Funda TURAN, Ayşegül ERGENLER & Ali UYAN**

Identification of a Potential Nursery Ground
of the Spiny Butterfly Ray, *Gymnura altavela*,
in the Northeastern Mediterranean Sea, Türkiye 205
Identifikacija potencialnih jasic za
metuljastega skata, Gymnura
altavela, v severovzhodnem
Sredozemskem morju,
Turčija

**Alen SOLDO, Eleonora DE SABATA &
Simona CLO**

Recent Records of Critically Endangered
Common Guitarfish *Rhinobatos rhinobatos*
(Linnaeus, 1758) in the Northern
Mediterranean 213
Nedavni zapiski o pojavljanju kritično
ogroženega navadnega goslaša
Rhinobatos rhinobatos
(Linnaeus, 1758) v
severnem Sredozemlju

Hakan KABASAKAL & F. Saadet KARAKULAK

Demersal Elasmobranchs of the Sea of Marmara:
Updated Inventory, Taxonomic Issues and
Environmental Implications 221
Pridnene hrustančnice v Marmarskem morju:
posodobljena inventarizacija, taksonomska
vprašanja in okoljske posledice

IHTIOLOGIJA

ITTILOGIA

ICHTHYOLOGY

**Jamila RIZGALLA, Amani FITORI, Francesco
TIRALONGO & Abdalrh BEN ABDALAH**

First Records of Blennies (Suborder
Blennioidea) off the Coast of Libya 245
Prvi zapisi o pojavljanju babic
(podred Blennioidea) ob
libijski obali

**Farid HEMIDA, Sara LADOUL, Christian
REYNAUD & Christian CAPAPÉ**

Confirmed Occurrence of the Mediterranean
Spearfish *Tetrapturus belone* (Osteichthyes:
Istiophoridae) from the Algerian Coast
(Southwestern Mediterranean Sea) 257
Potrjeno pojavljanje sredozemske
jadrovnice, Tetrapturus belone
(Osteichthyes: Istiophoridae) iz
alžirske obale (jugovzhodno
Sredozemsko morje)

Lana KHREMA, Adib SAAD & Christian CAPAPÉ First Substantiated Record of Blackfish <i>Centrolophus niger</i> (Centrolophidae) from the Syrian Coast (Eastern Mediterranean Sea) 263 <i>Prvi potrjen zapis črnuha, Centrolophus niger</i> (<i>Centrolophidae</i>), s sirske obale (vzhodno Sredozemsko morje)	MORSKA FAVNA <i>FAUNA MARINA</i> <i>MARINE FAUNA</i>
Amir IBRAHIM, Chirine HUSSEIN, Firas ALSHAWY & Alaa ALCHEIKH AHMAD Marine Fishes (Teleostei / Osteichthyes) of Syria (Eastern Mediterranean): an Updated Checklist 269 <i>Morske ribe (Teleostei / Osteichthyes)</i> <i>Sirije (vzhodno Sredozemsko morje):</i> <i>posodobljeni seznam vrst</i>	Nicola BETTOSO, Lisa FARESI, Saul CIRIACO, Marco FANTIN & Marco SEGARICH New Findings of the Cup-Shaped Demosponge <i>Calyx nicaensis</i> (Risso, 1826) on the Rocky Outcrops in the Gulf of Trieste (Northern Adriatic Sea) 311 <i>Nove najdbe morskega keliha Calyx nicaensis</i> (<i>Risso, 1826</i>) na skalnih osamelcih v Tržaškem zalivu (<i>severno Jadransko morje</i>)
BIOTSKA GLOBALIZACIJA <i>GLOBALIZZAZIONE BIOTICA</i> <i>BIOTIC GLOBALIZATION</i>	Aleksandra V. BORODINA & Yuri O. VELYAEV Features of Lipid Accumulation in Striped Venus Clam <i>Chamelea gallina</i> in the Sublittoral Zone of the Crimean Coast (Black Sea) 319 <i>Značilnosti kopičenja lipidov v navadni</i> <i>venerici (Chamelea gallina) v obrežnem</i> <i>pasu krimske obale (Črno morje)</i>
Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO, Enrico LA SPINA, Daniele TIBULLO & Francesco TIRALONGO <i>Cassiopea andromeda</i> at the Southernmost Tip of Italy: a Recent Arrival or an Overlooked Resident? 287 <i>Tujerodni klobučnjak Cassiopea andromeda</i> <i>na najjužnejši konici Italije: nedavni</i> <i>prišlek ali spregledan prebivalec?</i>	Okan AKYOL & Oğuzhan TAKICAK Recent Observations on <i>Monachus monachus</i> (Phocidae) at Sea-Cage Fish Farms in Izmir (Turkish Aegean Sea) 329 <i>Nedavna opažanja primerkov Monachus monachus</i> (<i>Phocidae</i>) v ribogojnicah z morskimi kletkami v <i>Izmirju (turško Egejsko morje)</i>
Mourad CHÉRIF, Rimel BENMESSAOUD, Sihem RAFRAFI-NOUIRA, Mohamed Nouri AYADI & Christian CAPAPÉ First Substantiated Record of the Golden-Banded Goatfish <i>Upeneus moluccensis</i> (Osteichthyes: Mullidae) from the Coast of Tunisia (Central Mediterranean Sea) 295 <i>Prvi potrjen zapis o zlatoprogem bradaču</i> <i>Upeneus moluccensis (Osteichthyes: Mullidae)</i> <i>z obale Tunizije (osrednje Sredozemsko morje)</i>	FLORA <i>FLORA</i> <i>FLORA</i>
Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ĆETKOVIĆ, Borut MAVRIČ & Lovrenc LIPEJ New Record of the Long-Jawed Squirrelfish, <i>Holocentrus adscensionis</i> (Osbeck, 1765), in the Adriatic Sea 303 <i>Novi zapis o pojavljanju veвериčjaka vrste</i> <i>Holocentrus adscensionis (Osbeck, 1765)</i> <i>v Jadranskem morju</i>	Amelio PEZZETTA & Marco PAOLUCCI La Flora di Palena (Parco Nazionale della Majella): Aggiornamento Floristico 337 <i>Flora Palene (Nacionalni park Majella):</i> <i>floristična posodobitev</i>
	DELO NAŠIH ZAVODOV IN DRUŠTEV <i>ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ</i> <i>ACTIVITIES BY OUR INSTITUTIONS AND</i> <i>ASSOCIATIONS</i>
	Bojana LIPEJ Reviving Landscapes, Connecting Species: Lessons from the ReCo Project 383
	IN MEMORIAM
	Tom TURK <i>V spomin Marjanu Richterju (1935–2025) 389</i>
	Kazalo k slikam na ovitku 391 <i>Index to images on the cover</i> 391

received: 2025-10-07

DOI 10.19233/ASHN.2025.32

NEW RECORD OF THE LONG-JAWED SQUIRRELFISH, *HOLOCENTRUS ADSCENSIONIS* (OSBECK, 1765), IN THE ADRIATIC SEA

Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ĆETKOVIĆ
Institute of Marine Biology, I Bokeljske brigade 68, 85330 Kotor, Montenegro

Borut MAVRIČ & Lovrenc LIPEJ
Marine Biology Station Piran, National Institute of Biology, Fornače 41, Slovenia
e-mail: Lovrenc.Lipej@nib.si

ABSTRACT

*On 16 October 2025, a specimen of the long-jawed squirrelfish, *Holocentrus adscensionis* (Osbeck, 1765), was caught in the waters of Porto Montenegro, Tivat (Boka Kotorska, Montenegro). This finding represents the first record of this non-indigenous species in the inventory of Montenegrin marine fish fauna and the second record in the Adriatic Sea.*

Key words: *Holocentrus adscensionis*, Holocentridae, bioinvasion, Boka Kotorska, Adriatic Sea

NUOVO AVVISTAMENTO DEL PESCE SCOIATTOLO, *HOLOCENTRUS ADSCENSIONIS* (OSBECK, 1765), NEL MAR ADRIATICO

SINTESI

*Il 16 ottobre 2025, un esemplare di pesce scoiattolo, *Holocentrus adscensionis* (Osbeck, 1765), è stato catturato nelle acque di Porto Montenegro, Tivat (Boka Kotorska, Montenegro). Questa scoperta rappresenta la prima registrazione di questa specie non autoctona nell'inventario della fauna ittica marina montenegrina e la seconda registrazione nell'Adriatico.*

Parole chiave: *Holocentrus adscensionis*, Holocentridae, bioinvasione, Boka Kotorska, Adriatico

INTRODUCTION

Climate change has been recognised as a major factor influencing Mediterranean biodiversity, alongside Atlantic influx, Lessepsian migration, and the introduction of exotic species by humans (Turan *et al.*, 2016). A similar trend is evident in the Adriatic Sea, where rapid biodiversity changes are occurring due to the increasing arrival of non-indigenous fishes and other taxa (Dulčić *et al.*, 1999; Lipej & Dulčić, 2004). The number of established alien species is expected to increase further in the near future, largely through the natural spread of populations already established in the central Mediterranean (Zenetos, 2010).

This paper presents a record of a non-indigenous fish species caught in the study area and explores possible explanations for its occurrence.

MATERIAL AND METHODS

Boka Kotorska is an 87 km² fjord-like bay in the eastern part of the southern Adriatic Sea, with a

coastline of 105.7 km. It comprises three distinct areas – the outer bay (Herceg Novi Bay), the middle bay (Tivat Bay), and the inner bay (Kotor-Risan Bay) – and is characterised by unique hydrological, geomorphological, climatological, and biotic factors (Gamulin-Brida, 1983) (Fig. 1). On 16 October 2025, Zoran Ćuk, a local fisherman, captured a specimen of an unknown fish while stick-float fishing at night near the Porto Montenegro marina (42.431497° N and 18.691224° E) in Tivat, located in the middle bay (Boka Kotorska, Montenegro). The catch occurred at a depth of approximately 11 m over a muddy bottom. The fisherman photographed the specimen – which had a total length of approximately 24 cm – and released it alive. The photograph was subsequently provided to the authors for analysis.

RESULTS AND DISCUSSION

The specimen (Fig. 2) exhibited all the typical characteristics that allowed it to be identified as *Holocentrus adscensionis* (Osbeck, 1765): body



Fig. 1: Map showing the location (Porto Montenegro, Tivat Bay, Montenegro) where the specimen of the long-jawed squirrelfish, *Holocentrus adscensionis*, was caught.

Sl. 1: Zemljevid z označeno lokaliteto (lokaliteta Porto Montenegro, Tivatski zaliv, Črna gora), kjer je bil ujet primerek veвериčjaka *Holocentrus adscensionis*.



Fig. 2: Specimen of the long-jawed squirrelfish, *Holocentrus adscensionis*, caught on 16 October 2025 in the waters of Porto Montenegro (Tivat, Boka Kotorska, Montenegro) (photo: Z. Ćuk).

Sl. 2: Primerek veвериčjaka, *Holocentrus adscensionis*, ujet 16. oktobra 2025 v vodah marine Porto Montenegro (Tivat, Boka Kotorska, Črna gora) (foto: Z. Ćuk).

oblong and laterally compressed, reddish with alternating red and white horizontal stripes; head pointed, with large eyes; upper jaw longer than the lower and extending beyond the centre of the eye; area between preopercle and opercle bearing rows of serrations; dorsal fin lacking the white spots behind the spine tips that are characteristic of the congeneric species *H. rufus* (Greenfield, 2003). The photograph obtained from the fisherman clearly shows the typical striped coloration, the eleven dorsal spines of subequal length lacking the white spots, and the evident preopercular spine. The caudal lobes are elongate, with the upper lobe markedly longer than the lower (*sensu* Woods, 1955).

The species *H. adscensionis* is a reef-associated fish of the family Holocentridae, occurring in the western Atlantic ranges from North Carolina (USA) and Bermuda to Brazil (Woods & Greenfield, 1978), and in the eastern Atlantic from Gabon to Ascension Island (Ben-Tuvia, 1990). It inhabits depths from shallow tide pools to 180 m, most commonly

between 8 and 30 m (Wyatt, 1983). It is a typically nocturnal species, spending the day hiding in crevices and cavities, and emerging at night to feed, mainly on crabs and other small crustaceans (Greenfield, 1981). The Porto Montenegro is a large, modern marina; however, its piers, overgrown with rich epibenthic fauna, contain many crevices and cracks that offer potential shelter for squirrelfish. It is also possible that the species occurs in adjacent areas, but due to its nocturnal behaviour it could easily be overlooked.

Prior to this study, there were no records of *H. adscensionis* from Montenegrin waters. In the Mediterranean Sea, the species had previously been reported only twice: first by Vella *et al.* (2016) in Maltese waters, and later by Ciriaco *et al.* (2022) from the Gulf of Trieste in the Adriatic Sea. The arrival of *H. adscensionis* is likely related to the gradual warming of the Mediterranean Sea (Occhipinti-Ambrogi & Galil, 2010). Novel arrivals via Atlantic influx are relatively uncommon in

the Adriatic Sea, since the main recognised entry route for non-indigenous fishes is the Suez Canal, through which more than 100 immigrant species have entered the Mediterranean Sea (Golani *et al.*, 2021). Vella *et al.* (2016) also considered the possibility that this species might have been introduced into Maltese waters through maritime activities in major ports. Given that *H. adscensionis* typically inhabits tropical and subtropical waters with temperatures ranging from approximately 23°C to 30°C (Reef Life Survey, 2025), we assume that winter water temperatures in Boka Kotorska are too low for the species to survive and become established.

To date, only records of seven alien fish species have been published in Montenegrin waters – including *Pterois miles* (Tomanić *et al.*, 2022) – which is relatively few compared to other Mediterranean regions. Based on the three occasional findings of the long-jawed squirrelfish in geographically disparate areas of the Mediterranean Sea (Malta, the Gulf of Trieste, and Boka Kotorska), it is too early to speculate on its spread or potential establishment in the area. However, the experience with its close relative, the redcoat, *Sargocentron rubrum* (Forsskål, 1775), first detected in the basin eighty years ago off Palestine marine waters ((Haas & Steinitz, 1947) and now considered one of the most successful colonisers in the Mediterranean Sea (Azzurro

et al., 2014) demonstrates that some species can rapidly find a niche in a new environment, establish themselves, and begin to spread. Furthermore, two new squirrelfish species have been reported for the first time in the Mediterranean Sea in recent years, namely *Neoniphon sammara* (Forsskål, 1775) (Deef, 2021; Mehanna & Osman, 2022) from Egyptian waters and *S. caudimaculatum* (Rüppell, 1838) from Tunisian waters (Ghanem *et al.*, 2022).

Although the specimen described here was not preserved in a formal museum collection, as recommended by best-practice guidelines (*sensu* Bello *et al.*, 2014), high-quality photographic evidence provided reliable documentation for confirming the species' presence (Dulčić *et al.*, 2006; Kovačić *et al.*, 2020). This case underscores the important role of citizen science in contributing data on alien species.

ACKNOWLEDGMENTS

The authors thank fisherman Zoran Ćuk for providing the photographic evidence and Nikša Miljanić for sharing the sighting of the squirrelfish. They also extend their gratitude to Tihomir Makovec and Branislav Lazarević for their support. The paper is a result of a bilateral scientific project between the Republic of Montenegro and the Republic of Slovenia (ARIS-BI/2024/205).

NOVI ZAPIS O POJAVLJANJU VEVERIČJAKA VRSTE *HOLOCENTRUS ADSCENSIONIS*
(OSBECK, 1765) V JADRANSKEM MORJU

Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ČETKOVIĆ
Institute of Marine Biology, I Bokaljske brigade 68, 85330 Kotor, Montenegro

Borut MAVRIČ & Lovrenc LIPEJ
Marine Biology Station Piran, National Institute of Biology, Fornače 41, Slovenia
e-mail: Lovrenc.Lipej@nib.si

POVZETEK

Ribič je 16. oktobra 2025 ujel primerek tujerodnega veveričjaka vrste *Holocentrus adscensionis* (Osbeck, 1765) v vodah marine Porto Montenegro (Tivat, Boka Kotorska, Črna gora). Gre za prvi zapis o pojavljanju te vrste za črnogorsko morsko ribjo favno in drugi zapis o pojavljanju v Jadranu.

Ključne besede: *Holocentrus adscensionis*, Holocentridae, bioinvazija, Boka Kotorska, Jadransko morje

REFERENCES

- Azzurro E., V.M. Tuset, A. Lombarte, F. Maynou, D. Simberloff, A. Rodríguez-Pérez & R.V. Solé (2014)**: External Morphology Explains the Success of Biological Invasions. *Ecol. Lett.*, 17, 1455-1463.
- Bello, G., R. Causse, L. Lipej & J. Dulčić (2014)**: A proposed best practice approach to overcome unverified and unverifiable “first records” in ichthyology. *Cybium*, 38(1), 9-14.
- Ben-Tuvia, A. (1990)**: Holocentridae. In: J.C. Quéro, J.C. Hureau, C. Karrer, A. Post and L. Saldanha (eds.): Check-list of the fishes of the eastern tropical Atlantic (CLOFETA), Vol. 2: pp. 627- 628. JNICT, Lisbon; SEI and UNESCO, Paris.
- Ciriaco, S., M. Segarich, V. Cirinà & L. Lipej (2022)**: First record of the Long-Jawed Squirrelfish *Holocentrus adscensionis* (Osbeck, 1765) in the Adriatic Sea. *Annales, Ser. Hist. Nat.*, 32(2), 309-316.
- Deef, L.E.M. (2021)**: First Record of Two Squirrelfishes, *Sargocentron spinosissimum* and *Sargocentron tiereoides* (Actinopterygii, Beryciformes, Holocentridae) From the Egyptian Mediterranean Coast. *Acta Ichthyol Piscat.*, 51(1), 107-112.
- Dulčić, J., B. Grbec & L. Lipej (1999)**: Information on the Adriatic ichthyofauna-effect of water warming? *Acta Adriatica*, 40, 33-43.
- Dulčić, J., B. Furlan & L. Lipej (2006)**: First confirmed record of *Lapanella fasciata* (Cocco, 1833) in the Adriatic Sea. *J. Applied Ichthyol.*, 22, 536-537.
- Gamulin-Brida, H. (1983)**: Crnogorsko primorje–specifični dio Jadrana s gledišta bentoskih biocenoza i njihove zaštite. *Studia Marina*, 13-14, 205-214.
- Ghanem, R., W. Bahri, A. Chaffai, J. Zaouali, B. Hassen, S. Karray, M. El Bour, J. Souissi & E. Azzurro (2022)**: First Record of the Silverspot Squirrelfish *Sargocentron caudimaculatum* (Rüppell, 1838) in Mediterranean Waters. *Front. Mar. Sci.*, 9, 869138. doi: 10.3389/fmars.2022.869138
- Greenfield, D.W. (1981)**: Holocentridae. In: W. Fischer, G. Bianchian, W.B. Scott (eds.) FAO species identification sheets for fishery purposes. Eastern Central Atlantic; fishing areas 34, 47. Department of Fisheries and Oceans Canada and FAO, 2.
- Golani, D., E. Azzurro, J. Dulčić, E. Massutí, & L. Orsi Relini (2021)**: Atlas of exotic fishes in the Mediterranean Sea (2nd edition). CIESM Publishers, 365 pp.
- Haas, G. & H. Steinitz (1947)**: Erythrean fishes on the Mediterranean coast of Palestine. *Nature*, 160, 28.
- Kovačić, M., L. Lipej & J. Dulčić (2020)**: Evidence approach to checklists: critical revision of the checklist of the Adriatic Sea fishes. *Zootaxa*, 4767(1), 1-55. <https://doi.org/10.11646/zootaxa.4767.1.1>.
- Lipej, L. & J. Dulčić (2004)**: The current status of Adriatic fish biodiversity. In: *Balkan Biodiversity: Pattern and Process in the European Hotspot*; Griffiths, H.I., Kryštufek, B., Reed, J.M., Eds.; Kluwer Academic: Dordrecht, The Netherlands; London, UK, 2004; pp. 291-306.
- Mehanna, S.F. & Y.A. Osman (2022)**: First record of the Lessepsian Sannara Squirrelfish, *Neoniphon sammara* (Forsskål, 1775), in the Egyptian Mediterranean waters. *Medit. Mar. Sci.*, 23(3), 664-667.
- Occhipinti Ambrogi, A. & B. Galil (2010)**: Marine alien species as an aspect of global change. *Advances in Oceanography and Limnology*, 1, 199-218. doi: 10.1080/19475721003743876.
- Reef Life Survey (2025)**: assessed on 30.10.2025 (www.reeflifesurvey.com)
- Tomanić, J., I. Četković, P. Simonović, Z. Ikica, A. Joksimović & A. Pešić (2022)**: New species of fish and crustaceans in Montenegrin waters (South Adriatic Sea). *Acta Adriatica*, 63, 109-122.
- Turan, C., D. Ergüden & M. Gürlek (2016)**: Climate Change and Biodiversity Effects in Turkish Seas. *Natural and Engineering Sciences*, 1, 15-24.
- Vella, A., N. Vella & S.A. Darmanin (2016)**: The first record of the longjaw squirrelfish, *Holocentrus adscensionis* (Osbeck, 1765) (Holocentridae: Holocentridae), in the Mediterranean Sea. *Natural and Engineering Sciences*, 1(3), 78-85. doi: 10.28978/nesciences.286371.
- Woods, L.P. (1955)**: Western Atlantic species of the genus *Holocentrus*. *Fieldiana. Zoology*, 37(4), 91-119.
- Woods, L.P. & D.W. Greenfield (1978)**: Holocentridae. In: W. Fischer (ed.) FAO species identification sheets for fishery purposes. Western Central Atlantic (Fishing Area 31). Vol. 3. FAO, Rome.
- Wyatt, J.R. (1983)**: The biology, ecology and bionomics of the squirrelfishes, Holocentridae. In: J.L. Munro (ed.) Caribbean coral reef fishery resources. *ICLARM Stud.*, 7, 50-58.
- Zenetos, A. (2010)**: Trend in Alien species in the Mediterranean. An answer to Galil, 2009 ‘Taking stock: inventory of alien species in the Mediterranean Sea. *Biol Invasions*, 12, 3379-3381.