

# ANNALES

*Anali za istrske in mediteranske študije*  
*Annali di Studi istriani e mediterranee*  
*Annals for Istrian and Mediterranean Studies*  
*Series Historia Naturalis, 36, 2026, 1*





# ANNALES

**Anali za istrske in mediteranske študije  
Annali di Studi istriani e mediterraneei  
Annals for Istrian and Mediterranean Studies**

**Series Historia Naturalis, 36, 2026, 1**

**UREDNIŠKI ODBOR/  
COMITATO DI REDAZIONE/  
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Ana Fortič, Andrej Gogala, Daniel Golani (IL), Danijel Ivajnsič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačić (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, James Pierson (USA), Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/  
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/  
Redattore responsabile per le scienze  
naturali/Natural Science Editor:**

Lovrenc Lipej

**Urednici/Redattrici/Editors:**

Martina Orlando-Bonaca, Ana Fortič

**Prevajalci/Traduttori/Translators:**

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/  
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

**Tisk/Stampa/Print:**

Založništvo PADRE d.o.o.

**Izdajatelj/Editori/Published by:**Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®  
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/  
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran  
SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;  
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2026.

**Sofinancirajo/Supporto finanziario/  
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

**Annales - Series Historia Naturalis** izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>  
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>  
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / *CONTENTS*

BIOTSKA GLOBALIZACIJA  
*GLOBALIZZAZIONE BIOTICA*  
*BIOTIC GLOBALIZATION*

**Alen SOLDO, Rigers BAKIU & Sherif DURMISHAJ**

The First Record of Bastard Grunt *Pomadasys incisus* (Bowdich, 1825) in Albanian Waters (Adriatic Sea) ..... 1  
*Prvi zapis o pojavljanju neprave prašičevke Pomadasys incisus (Bowdich, 1825) v albanskih vodah (Jadransko morje)*

**Pero UGARKOVIĆ, Ilija ČETKOVIĆ, Olivera MARKOVIĆ, Aleksandar JOKSIMOVIĆ, Nikola ĐORĐEVIĆ & Jakov DULČIĆ**

Additional Records of the Bastard Grunt, *Pomadasys incisus* (Bowdich, 1825), from the Eastern Adriatic Sea ..... 7  
*Dodatne najdbe neprave prašičevke, Pomadasys incisus (Bowdich, 1825), iz vzhodnega Jadranskega morja*

**Rüştü KIRMAN & Murat BİLECENOĞLU**

Significant Range Expansion of *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) in the Aegean Sea Based on Scuba and Photographic Observations ..... 15  
*Znatno razširjanje areala vrste Sepioteuthis lessoniana (Cephalopoda: Loliginidae), v Egejskem morju na podlagi podvodnih opazovanj in fotografiranja*

**Adrian BRAJKOVIĆ, Iris MATULJA & Neven IVEŠA**

Conquering the North: New Occurrence of the Common Lionfish, *Pterois miles* (Bennett, 1828), in the Adriatic Sea ..... 21  
*Osvajanje severa: novi zapis o pojavljanju navadne plamenke, Pterois miles (Bennett, 1828), v Jadranskem morju*

**Alan DEIDUN, Bruno ZAVA, Maria CORSINI-FOKA, Arnold SCIBERRAS & Alessio MARRONE**

First Record of the Devil Firefish *Pterois miles* (Bennett, 1828) (Actinopterygii: Scorpaenidae) from Maltese Coastal Waters ..... 29  
*Prvi zapis o pojavljanju plamenke Pterois miles (Bennett, 1828) (Actinopterygii: Scorpaenidae) v malteških obalnih vodah*

**Chirine HUSSEIN, Amir IBRAHIM, Firas ALSHAWY, Mouina BADRAN & Rahaf ABO ASA**

First Mediterranean Record of the Diagonal Butterflyfish, *Chaetodon fasciatus* Forsskål, 1775, Reported from Syrian Waters ..... 35  
*Prvi sredozemski zapis o pojavljanju poševnoprogame ščetinozobke, Chaetodon fasciatus Forsskål, 1775, iz sirske vode*

**Houssein ELBARAASI, Tarek SHOEIB, Mona SAID & Laith A. JAWAD**

A Further Record of the Barred Knifejaw, *Oplegnathus fasciatus* (Temminck & Schlegel, 1844), a Pacific Species, from the Mediterranean: A New Record from Benghazi, Libya ..... 41  
*Novi zapis o vrsti Oplegnathus fasciatus (Temminck & Schlegel, 1844), pacifiški vrsti, v Sredozemlju: nova najdba iz Bengazija v Libiji*

**Alen SOLDO & Rigers BAKIU**

Rapid Increase in Records of the Invasive Silver-cheeked Toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Adriatic Sea ..... 49  
*Hitro naraščanje pojavov invazivne srebrnoproge napihvalke Lagocephalus sceleratus (Gmelin, 1789) v Jadranskem morju*

**Gerasimos KONDYLATOS, Konstantinos KALAENTZIS, Styliani MINOUDI & Maria CORSINI-FOKA**

Further Molecular Identification Confirms the Occurrence of *Lagocephalus guentheri* Miranda Ribeiro, 1915 in the Aegean Coastal Waters of Greece ..... 55  
*Nadaljnja molekularna identifikacija potrjuje prisotnost vrste Lagocephalus guentheri Miranda Ribeiro, 1915 v egejskih obalnih vodah Grčije*

**Deniz ERGÜDEN & Cem ÇEVİK**

Length–Weight and Length–Length Relationships, and Condition Factor of *Ambassis dussumieri* Cuvier, 1828, in the Northeastern Mediterranean, Türkiye ..... 65  
*Dolžinsko-masni odnos med dolžino in kondicijskim faktorjem pri vrsti Ambassis dussumieri Cuvier, 1828, v severovzhodnem Sredozemskem morju, Turčija*

**Okun AKYOL & HALİL ŞEN**

Unexpected Occurrence of Prussian Carp *Carassius gibelio* (Cyprinidae) in Homa Lagoon (Izmir Bay, Aegean Sea) ..... 73  
*Nepričakovani pojav srebrnega koreslja Carassius gibelio (Cyprinidae) v laguni Homa (Izmirski zaliv, Egejsko morje)*

SREDOZEMSKA HRUSTANČNICE  
 SQUALI E RAZZE MEDITERRANEE  
 MEDITERRANEAN SHARKS AND RAYS

**Terry CARBON, Emily GIGNON, Justine LALLAU-VAZZOLER, Hugo MENARD, Claudio BARRIA, Ana I. COLMENERO & Nicolas ZIANI**

Precopulatory Behaviour of *Pteroplatytrygon violacea* (Myliobatiformes: Dasyatidae) in the Northwestern Mediterranean ..... 81  
*Predkopulacijsko vedenje vijoličnega morskega biča (Pteroplatytrygon violacea) (Myliobatiformes: Dasyatidae) v severozahodnem Sredozemlju*

<b>Hakan KABASAKAL</b> Distribution of <i>Odontaspis ferox</i> in the Mediterranean Sea: Insights from Spatial and Temporal Analyses ..... 87 <i>Razširjenost vrste Odontaspis ferox v Sredozemskem morju: Spoznanja na podlagi prostorskih in časovnih analiz</i>	MORSKA FAVNA FAUNA MARINA MARINE FAUNA
<b>Hristina GELEVSKA, Borut MAVRIČ, Lovrenc LIPEJ &amp; Christian CAPAPÉ</b> Is the Gulf of Trieste a potential nursery area for some elasmobranch species? ..... 99 <i>Ali je Tržaški zaliv potencialno območje odraščanja (jaslice) za določene vrste hrustančnic?</i>	<b>Francesco TIRALONGO, Paola LEOTTA &amp; Riccardo MARTELLUCCI</b> <i>Physalia physalis</i> in the central Mediterranean Sea: Recent observations associated with mass strandings of <i>Velella velella</i> and surface circulation dynamics ..... 165 <i>Portugalska ladjica (Physalia physalis) v osrednjem Sredozemskem morju: nedavna opažanja v povezavi z masovnimi nasedanji morskega jadrčka (Velella velella) in dinamiko površinskega kroženja vode</i>
IHTIOFAVNA ITTIOFAUNA ICHTHYOFAUNA	
<b>Cem DALYAN, Yunus GÖNÜL, Mahmud Samed ŞAHİNOĞULLARI &amp; Hakan KABASAKAL</b> New data on the Occurrence and Morphology of the Armless Snake Eel, <i>Dalophis imberbis</i> (Ophichthidae), from the Northeastern Mediterranean Sea ..... 119 <i>Novi podatki o pojavljanju in morfologiji kačaste jegulje, Dalophis imberbis (Ophichthidae), iz severovzhodnega Sredozemskega morja</i>	<b>Andrea LOMBARDO, Juba CHABANE &amp; Rachida GHALMI</b> First Contribution to the Study of the "Sea Slug" Fauna (Gastropoda, Heterobranchia) of the Algerian coast ..... 177 <i>Prvi prispevek k proučevanju favne „goliš morskih polžev“ (Gastropoda, Heterobranchia) alžirske obale</i>
<b>Chirine HUSSEIN, Firas ALSHAWY &amp; Amir IBRAHIM</b> First Record of the Mediterranean Dealfish, <i>Trachipterus trachipterus</i> (Gmelin, 1789) (Trachipteridae), in Syrian Marine Waters ..... 127 <i>Prvi zapis o pojavljanju kosice, Trachipterus trachipterus (Gmelin, 1789) (Trachipteridae), v sirskih morskih vodah</i>	<b>Borut MAVRIČ, Tjaša PREMRL, Martin MAVRIČ, Tina MIRT, Neža LEBAN, Tihomir MAKOVEC &amp; Lovrenc LIPEJ</b> Assessing Soft-Bottom Epibenthic Communities: Methodological Insights into Dredging and Video Surveys ..... 203 <i>Ocenjevanje epibentoških združb mehkega dna: metodološki vpogledi v dredžanje in video pregleda</i>
<b>Igor AGOSTINI &amp; Okan AKYOL</b> On the Occurrence of the Currently Largest Recorded <i>Pagellus erythrinus</i> (Sparidae) in the Tyrrhenian Sea (Italy) ..... 133 <i>O pojavljanju trenutno največjega evidentiranega primerka vrste Pagellus erythrinus (Sparidae) v Tirenskem morju (Italija)</i>	FAVNA FAUNA FAUNA
<b>Ismail Burak DABAN &amp; Yusuf ŞEN</b> Substantiated Record and Preliminary Biological Insights of <i>Microlipophrys dalmatinus</i> (Steindachner & Kolombatović, 1883) from the Sea of Marmara, Türkiye ..... 141 <i>Potrjen zapis o pojavljanju in predhodna biološka spoznanja o vrsti Microlipophrys dalmatinus (Steindachner &amp; Kolombatović, 1883) iz Marmarskega morja, Turčija</i>	<b>Kim LEBAN, Iztok ŠKORNIK, Špela ČONČ, Mateja BREG VALJAVEC, Žan KURALT, Lenart ŠTAUT, Jure TIČAR, Katarina POLAJNAR HORVAT, Primož GAŠPERIČ, Aleš SMREKAR &amp; Melita VAMBERGER</b> Habitat Insights for the European Pond Turtle ( <i>Emys orbicularis</i> ) in the Coastal Wetland Area Sečovlje Salina Nature Park, Slovenia ..... 221 <i>Habitatne značilnosti močvirske sklednice (Emys orbicularis) v krajinskem parku Sečoveljske soline, Slovenija</i>
<b>Zeliha ERDOĞAN, Gülçin ULUNEHİR AYDIN, Hatice TORCU-KOÇ &amp; Tuğba ANBAROĞLU</b> Population Study of <i>Trachurus mediterraneus</i> Focused on Reproductive Biology in Edremit Bay, Northern Aegean Sea ..... 151 <i>Razmnoževalna biologija sredozemskega šura (Trachurus mediterraneus) v sklopu populacijske raziskave v Edremitnem zalivu (severno Egejsko morje)</i>	<b>Leon CIMERMAN &amp; Matija KRIŽNAR</b> Najdbe pleistocenske avifavne v kamnolomu Črni Kal (Primorska, Slovenija) ..... 243 <i>Pleistocene Avifauna Finds in the Črni Kal Quarry (Primorska Region, Slovenia)</i>
	Kazalo k slikam na ovitku ..... 255 <b>Index to images on the cover</b> ..... 255

## THE FIRST RECORD OF BASTARD GRUNT *POMADASYS INCISUS* (BOWDICH, 1825) IN ALBANIAN WATERS (ADRIATIC SEA)

*Alen SOLDO*

Faculty of Marine Sciences, University of Split, Croatia  
e-mail: soldo@unist.com

*Rigers BAKIU*

Department of Aquaculture and Fisheries, Faculty of Agriculture and Environment, Agricultural University of Tirana, Koder-Kamez,  
Albania

ACEPSD, Albanian Center for Environmental Protection and Sustainable Development, Tirana, Albania

*Sherif DURMISHAJ*

Qendra e Peshkimit Oriku, Radhime, Vlore, Albania

### ABSTRACT

The bastard grunt *Pomadasys incisus* (Bowdich, 1825), an Atlantic species that entered the Mediterranean through the Strait of Gibraltar, exhibits a discontinuous distribution within the basin and remains rare in the Adriatic Sea. Here, we report the first record of *P. incisus* in Albanian waters and only the third confirmed occurrence in the Adriatic Sea. A single individual was captured on 9 October 2025 in the Bay of Vlorë at a depth of 48 m near aquaculture facilities. The specimen was identified based on diagnostic morphological features and measured 21 cm in total length and 125 g in weight. This record fills an important gap in the species' Mediterranean distribution, although the exact route of entry into the Adriatic remains unclear. The rarity of records suggests limited establishment; however, recent warming trends in the Adriatic may facilitate further expansion.

**Key words:** *Pomadasys incisus*, bastard grunt, Albanian Coast, Adriatic Sea, Mediterranean Sea

## PRIMO RITROVAMENTO DI *POMADASYS INCISUS* (BOWDICH, 1825) NELLE ACQUE ALBANESI (MARE ADRIATICO)

### SINTESI

Il grugnitore bastardo *Pomadasys incisus* (Bowdich, 1825), una specie atlantica entrata nel Mediterraneo attraverso lo Stretto di Gibilterra, presenta una distribuzione discontinua all'interno del bacino e rimane raro nell'Adriatico. Gli autori riportano il primo ritrovamento di *P. incisus* nelle acque albanesi e solo la terza presenza confermata nell'Adriatico. Un singolo individuo è stato catturato il 9 ottobre 2025 nella Baia di Valona a una profondità di 48 m, in prossimità di impianti di acquacoltura. L'esemplare è stato identificato sulla base di caratteristiche morfologiche diagnostiche e misurava 21 cm di lunghezza totale e 125 g di peso. La rarità delle segnalazioni suggerisce un insediamento limitato; tuttavia, i recenti trend di riscaldamento dell'Adriatico potrebbero favorire un'ulteriore espansione.

**Parole chiave:** *Pomadasys incisus*, grugnitore bastardo, costa albanese, Adriatico, Mediterraneo

## INTRODUCTION

The family Haemulidae comprises approximately 138 species distributed among 23 genera, inhabiting predominantly coastal waters of tropical and subtropical regions across the Atlantic, Indian, and Pacific Oceans. Members of this family are mainly marine, although some species occur in brackish environments and only a few inhabit freshwater systems (Fricke *et al.*, 2022; Van der Laan *et al.*, 2022). Within this family, the genus *Pomadasys* includes 29 species, of which three have been reported in the Mediterranean Sea: the striped piggy *Pomadasys stridens* (Forsskål, 1775), the bastard grunt *Pomadasys incisus* (Bowdich, 1825), and the silver grunt *Pomadasys argenteus* (Forsskål, 1775). Notably, all three species are considered non-indigenous to the Mediterranean basin.

Two of these species, *P. stridens* and *P. argenteus*, are Lessepsian migrants that entered the Mediterranean Sea from the Red Sea via the Suez Canal. *Pomadasys stridens*, native to the Indo-Pacific region, was first recorded in the Mediterranean around 1888 in Port Said and has since established populations throughout the eastern Mediterranean, where it is considered one of the most abundant Lessepsian fish, but also occasionally reaches the Italian coast (Golani *et al.*, 2002; Bilecenoglu *et al.*, 2009; Bodilis *et al.*, 2013; Tüzün & Gücü, 2023). More recently, *P. argenteus* was recorded for the first time in the Mediterranean in 2022 near Misrata, Libya, representing the second Red Sea haemulid species to colonize the basin (Fitori *et al.*, 2023).

In contrast, *Pomadasys incisus* has an Atlantic origin. Originally described from Gambia, it is widely distributed along the eastern Atlantic coastline, ranging from South Africa to the Iberian Peninsula, including Macaronesian archipelagos such as the Canary Islands, Madeira, and Cape Verde. The species entered the Mediterranean Sea through the Strait of Gibraltar, probably during the mid-19th century, likely facilitated by favorable oceanographic conditions, including prevailing currents, increasing sea temperatures, and the availability of suitable habitats, particularly soft substrates in shallow waters (Francour *et al.*, 1994; Bodilis *et al.*, 2013; Doumpas *et al.*, 2020; Kovačić *et al.*, 2021). Following its introduction, *P. incisus* established populations in the northwestern Mediterranean, where it has shown increasing abundance in areas such as the Málaga coast, the Catalan Sea, and the Gulf of Lion (Bodilis *et al.*, 2013; Villegas-Hernández *et al.*, 2014).

Despite its first documented occurrence in the Mediterranean dating back to 1840, it is possible that *P. incisus* was present earlier but remained undetected until the mid-19th century. Since then, the species has been recorded across a wide range of Mediter-

anean localities, including the coasts of Algeria, Tunisia, Spain, France, Italy, Libya, Egypt, Israel, Lebanon, Syria, Turkey, Greece, Malta, Cyprus and Croatia (Bodilis *et al.*, 2013; Doumpas *et al.*, 2020; Çoker & Akyol, 2021). Nevertheless, its distribution remains discontinuous and locally variable, suggesting a complex colonization pattern.

The successful dispersal of *P. incisus* is likely linked to its broad ecological tolerance and life-history traits. This gonochoric species exhibits relatively fast growth and a moderate lifespan of up to seven years (Pajuelo *et al.*, 2003a,b). It inhabits both marine and brackish environments, typically at depths ranging from 10 to 100 m, and occupies a wide variety of habitats, including sandy and rocky substrates, seagrass meadows, as well as lagoons and estuarine systems. Its feeding ecology is similarly opportunistic, consisting mainly of benthic invertebrates. Reproductive patterns appear to vary geographically, e.g., in warmer waters of the eastern Atlantic and the southern central Mediterranean, spawning takes place during a more prolonged season from June to November, while in the north-western Catalan Sea spawning season is shorter, occurring between July and September, and between August and October in the Gulf of Tunis, while, in contrast, all year round in the Canarian Archipelago (Pajuelo *et al.*, 2003a, 2003b; Fehri-Bedoui & Gharbi, 2008; Kapiris *et al.*, 2008; Bodilis *et al.*, 2013; Villegas-Hernández *et al.*, 2014; Doumpas *et al.*, 2020).

Despite this ecological flexibility and relatively broad distribution, *P. incisus* remains poorly studied in the Mediterranean, and available data on its biology and population dynamics are limited (Kapiris *et al.*, 2008). Moreover, its occurrence is still sporadic, with only a few areas reporting consistent or high local abundances. This pattern suggests that both intrinsic biological factors and extrinsic environmental constraints may influence its establishment success and limit its wider expansion across the Mediterranean basin (Bodilis *et al.*, 2013).

In this context, documenting new occurrences is essential for improving our understanding of the species' distribution and ongoing biogeographical changes in the Mediterranean Sea. Here, we report the first record of *Pomadasys incisus* in Albanian waters, representing only the third confirmed occurrence of the species in the Adriatic Sea. This finding contributes to filling a significant gap in the known distribution of the species and provides further insight into its range expansion within the Mediterranean basin.

## MATERIAL AND METHODS

On 9 October 2025, a single individual of *Pomadasys incisus* was accidentally captured by a trammel net in the Bay of Vlora (southern Albania)

(40.444600 °N, 19.383164 °E) (Fig. 1). The specimen was caught at a depth of 48 m, approximately 500 m from the nearest coastline, in proximity to marine aquaculture cages. The seabed in this area is predominantly muddy, likely influenced by organic enrichment associated with fish farming activities. After capture, the specimen was brought ashore, then photographed and measured (length to the nearest millimeter and weight in grams) by one of the authors (S. Durmishaj). Since the specimen was part of a commercial catch, it was returned to the fishermen and later sold at the Fishing Center Oriku in Radhimë, a coastal village near the city of Vlorë.



**Fig. 1:** The map of the area with the indicated location (\*) of the record of *Pomadasys incisus*.

**Sl. 1:** Zemljevid obravnavanega območja z označeno lokality ulova (\*) vrste *Pomadasys incisus*.

## RESULTS AND DISCUSSION

The specimen was identified as *Pomadasys incisus* based on its diagnostic morphological features, including a relatively large head with a depth comparable to that of the body, a small and slightly oblique mouth, greyish coloration on the dorsum, and silvery-white coloration on the flanks and ventral side. Additional distinguishing characteristics included a dark spot on the posterior margin of the operculum and orange to yellowish fins (Fig. 2). These traits are consistent with previous descriptions of the species (Kapiris *et al.* 2008; Bodilis *et al.*, 2013; Akyol & Ünal, 2016; Doumpas *et al.*, 2020). Hence, the individual measured 21.0 cm in total length (TL) and weighed 125 g.

This record represents the first confirmed occurrence of *P. incisus* in Albanian waters and only the third documented sighting of the species in the Adriatic Sea. The first Adriatic record dates back to 15 August 2015, when a single individual was captured off the Pelješac Peninsula (Croatia) in the southern Adriatic (Karachle *et al.*, 2016). The second known record was recently reported from the western side of the Adriatic Sea, when on 5 July 2025 a specimen was caught by a recreational fisher using rod and line in coastal waters at a depth of 25 m off Ancona harbour (Toma *et al.*, 2025). No additional records have been reported in the Adriatic basin until the present study, suggesting either a very low detection probability or a sporadic occurrence pattern in this region.

The present finding is consistent with previously proposed dispersal pathways of *P. incisus* within the Mediterranean Sea (Bodilis *et al.*, 2013). After entering through the Strait of Gibraltar, the species is thought to expand via two main routes: (1) an eastward pathway along the North African coastline, from Algeria to Tunisia and further into the Levantine basin, and (2) a northward pathway along the northern Mediterranean coasts, initially following Spain and France and subsequently extending toward other northern regions.

However, the occurrence of *P. incisus* in the Adriatic Sea raises questions regarding the exact dispersal route taken. Given the semi-enclosed nature of the Adriatic basin and its connection to the Mediterranean through the relatively narrow Strait of Otranto, individuals may have entered from the Ionian Sea, potentially originating from populations established along the North African coast (e.g., Gulf of Tunis). Alternatively, a west-to-east dispersal route via the Tyrrhenian Sea cannot be excluded.

It is also important to consider the environmental constraints of the Adriatic Sea, which is generally recognized as one of the coldest sub-basins of the Mediterranean. Such conditions may have historically



**Fig. 2:** The left (A) and right (B) sides of the caught *Pomadasys incisus*.  
**Sl. 2:** Leva (A) in desna stran (B) ujetega primerka vrste *Pomadasys incisus*.

limited the establishment and persistence of thermophilic species such as *P. incisus*. Nevertheless, recent oceanographic trends indicate a significant increase in sea temperature and salinity in the Adriatic over the past decade (Terzić *et al.*, 2025). These changes are likely facilitating the northward expansion of warm-affinity species, as increasingly documented across the Mediterranean basin.

In this context, the present record may reflect an ongoing shift in species distribution driven by climate-related changes. The sporadic but recurrent appearance of *P. incisus* in the Adriatic could represent early stages of colonization, although further records are needed to determine whether a stable population will establish.

Finally, such occurrences highlight the importance of reporting single records, as they can serve as early

indicators of ecological change. The detection of non-indigenous or range-expanding thermophilic species may provide valuable insights into shifting environmental conditions and habitat suitability, often preceding detectable trends in long-term oceanographic datasets.

#### ACKNOWLEDGMENTS

This research was funded by the European Union (NextGenerationEU) under the Croatian Recovery and Resilience Plan 2021–2026 (NRRP), through the University of Split institutional project “Anthropogenic Impact on the Sustainability of Coastal Marine Ecosystems - AIME; IP-UNIST-52”, approved by the Ministry of Science, Education and Youth of the Republic of Croatia.

## PRVI ZAPIS O POJAVLJANJU NEPRAVE PRAŠIČEVKE *POMADASYS INCISUS* (BOWDICH, 1825) V ALBANSKIH VODAH (JADRANSKO MORJE)

*Alen SOLDI*

Faculty of Marine Sciences, University of Split, Croatia  
e-mail: soldo@unist.com

Rigers BAKIU

Department of Aquaculture and Fisheries, Faculty of Agriculture and Environment, Agricultural University of Tirana, Koder-Kamez, Albania  
ACEPSD, Albanian Center for Environmental Protection and Sustainable Development, Tirana, Albania

*Sherif DURMISHAJ*

Qendra e Peshkimit Oriku, Radhime, Vlore, Albania

### POVZETEK

Neprava prašičevka, *Pomadasys incisus* (Bowdich, 1825), atlantska vrsta, ki je v Sredozemsko morje vstopila skozi Gibraltarsko ožino, kaže prekinjeno razširjenost znotraj bazena in ostaja redka v Jadranskem morju. Avtorji poročamo o prvi najdbi vrste *P. incisus* v albanskih vodah in šele o tretjem potrjenem pojavljanju v Jadranskem morju. Primerek te vrste je bil ulovljen 9. oktobra 2025 v Vlorovskem zalivu na globini 48 m v bližini marikulturnih objektov. Primerek smo določili na podlagi diagnostičnih morfoloških značilnosti ter je meril 21 cm v celotno dolžino in tehtal 125 g. Ta zapis o pojavljanju zapolnjuje pomembno vrzel v sredozemski razširjenosti te vrste, čeprav natančen način njenega vstopa v Jadransko morje ostaja nejasen. Redkost najdb kaže na omejeno ustalitev; vendar pa bi nedavni trendi segrevanja Jadranskega morja lahko olajšali nadaljnjo razširjanje.

**Ključne besede:** *Pomadasys incisus*, neprava prašičevka, albanska obala, Jadransko morje, Sredozemsko morje

## REFERENCES

- Akyol, O. & V. Ünal (2016):** First Record of a Lessepsian Migrant, *Pomadasys stridens* (Actinopterygii: Perciformes: Haemulidae), From the Aegean Sea, Turkey. *Acta Ichthyol. Piscat.*, 46(1), 53–55.
- Bilecenoglu, M., M. Kaya & A. Eryigit (2009):** New data on the occurrence of two alien fishes, *Pisodonophis semicinctus* and *Pomadasys stridens*, from the Eastern Mediterranean Sea. *Mediterr. Mar. Sci.*, 10(2), 151–155.
- Bodilis, P., F. Crocetta, J. Langeneck & P. Francour (2013):** The spread of an Atlantic fish species, *Pomadasys incisus* (Bowdich, 1825) (Osteichthyes: Haemulidae), within the Mediterranean Sea with new additional records from the French Mediterranean coast. *Ital. J. Zool.*, 80(2), 273–278.
- Çoker, T. & O. Akyol (2021):** On the occurrence of *Pomadasys incisus* (Haemulidae) in the Turkish Aegean Sea (Eastern Mediterranean Sea). *Annales, Ser. Hist. Nat.*, 31(1), 123–128.
- Doumpas, N., V. Tanduo, F. Crocetta, I. Giovos, J. Langeneck, F. Tiralongo & P. Kleitou (2020):** The bastard grunt *Pomadasys incisus* (Bowdich, 1825) (Teleostei: Haemulidae) in Cyprus (eastern Mediterranean Sea) – a late arrival or just a neglected species? *Biodivers. Data J.*, 8, e58646.
- Fehri-Bedoui, R. & H. Gharbi (2008):** Sex-ratio, reproduction and feeding habits of *Pomadasys incisus* (Haemulidae) in the Gulf of Tunis (Tunisia). *Acta Adriat.*, 49(1), 5–19.
- Fituri, A., A.E. Fituri, & D. Golani (2023):** The probable first record of the silver grunt, *Pomadasys argenteus* (Forsskål, 1775), from the Mediterranean Sea. *Cah. Biol. Mar.*, 64, 179–181.
- Francour, P., C. F. Boudouresque, J.G. Harmelin, M.L. Harmelin-Vivien & J.P. Quignard (1994):** Are the Mediterranean waters becoming warmer? Information from biological indicators. *Mar. Pollut. Bull.*, 28(9), 523–526.
- Fricke, R., W. N. Eschmeyer & R. Van der Laan (eds.) (2022):** Eschmeyer's catalogue of fishes: genera, species, references. California Academy of Sciences, San Francisco. Available online: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>
- Golani, D., L. Orsi-Relini, E. Massutí & J.P. Quignard, (2002):** CIESM atlas of exotic species in the Mediterranean. Vol. 1: Fishes. CIESM Publications, Monaco, 256 pp.
- Kapiris, K., E. Kallias, & A. Conides (2008):** Preliminary biological data on *Pomadasys incisus* (Osteichthyes: Haemulidae) in the Aegean Sea, Greece. *Mediterr. Mar. Sci.*, 9(2), 53–62.
- Karachle, P., A. Angelidis, G. Apostolopoulos, D. Ayas, M. Ballesteros, C. Bonnici, M. Brodersen, L. Castriota, N. Chalari, J. Cottalorda, F. Crocetta, A. Deidun, Z. Đodo, A. Dogrammatzi, J. Dulčić, F. Fiorentino, O. Gonulal, J. Harmelin, G. Insacco, D. Izquierdo-Gómez, A. Izquierdo-Muñoz, A. Joksimović, S. Kavadas, M. Malaquias, E. Madrenas, D. Massi, P. Micarelli, D. Minchin, U. Onal, P. Ovalis, D. Poursanidis, A. Siapatis, E. Sperone, A. Spinelli, C. Stamouli, F. Tiralongo, S. Tuncer, D. Yağlıoğlu, B. Zava & A. Zenetos (2016):** New Mediterranean biodiversity records (March 2016). *Mediterr. Mar. Sci.*, 17(1), 230–252.
- Kovačić, M., L. Lipej, J. Dulčić, S.P. Iglesias & M. Goren (2021):** Evidence-based checklist of the Mediterranean Sea fishes. *Zootaxa*, 4998(1), 1–115.
- Pajuelo, J.G., J.M. Lorenzo & M. Gregoire (2003a):** Age and growth of the bastard grunt (*Pomadasys incisus*) (Haemulidae) inhabiting the Canarian archipelago, Northwest Africa. *Fish. Bull.*, 101(4), 851–859.
- Pajuelo, J.G., J.M. Lorenzo, M. Gregoire & R. Domínguez-Seoane (2003b):** Life history of *Pomadasys incisus* (Osteichthyes: Haemulidae) of the Canarian Archipelago. *Sci. Mar.*, 67(2), 241–248.
- Terzić, E., V. Cardin, J. Le Meur, N. Dunić, M. Vodopivec & I. Vilibić (2025):** Unprecedented warming and salinization observed in the deep Adriatic. *Limnol. Oceanogr. Lett.*, 10, 888–898.
- Tüzün, S. & A.C. Gücü, (2023):** Diet variations of striped piggy, *Pomadasys stridens* (Forsskål, 1775) (Teleostei: Haemulidae) in the northeastern Mediterranean Sea. *Reg. Stud. Mar. Sci.*, 66, 103129.
- Toma, M., D. Agius, E. Azzurro, M. Bo, F. Crocetta, F., I. Dal, T. Delli Carri, M.C. Deval, L. Galli, A. Gkoulia, F. Grati, D. Kolokotronis, L. Lipej, C. Matthews, B. Mavrič, R. Mejri, J. Murcia, E.S. Okudan, V. Orenes-Salazar, J.P. Renoult, L. Pisani, J. Rizgalla, F. Russo, A. Salman, A. Soldo, V. Tanduo, F. Tiralongo, I. Tuney & A. Ünlüoğlu (2025):** New records of rarely reported species in the Mediterranean Sea (December 2025). *Mediterr. Mar. Sci.*, 26(4), 972–994.
- Van der Laan, R., R. Fricke & W.N. Eschmeyer (eds.) (2022):** Eschmeyer's catalogue of fishes: classification. California Academy of Sciences, San Francisco. Available online: <http://www.calacademy.org/scientists/catalog-of-fishes-classification/>
- Villegas-Hernández, H., J. Lloret & M. Muñoz (2014):** Climate-driven changes in life-history traits of the bastard grunt (*Pomadasys incisus*) in the north-western Mediterranean. *Mediterr. Mar. Sci.*, 16(1), 21–30.